

TREX2 Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP21082a

Specification

TREX2 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	<u>Q9BQ50</u>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	25922

TREX2 Antibody (C-term) - Additional Information

Gene ID 11219

Other Names Three prime repair exonuclease 2, 3'-5' exonuclease TREX2, TREX2

Target/Specificity

This TREX2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 264-299 amino acids from the C-terminal region of human TREX2.

Dilution WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

TREX2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

TREX2 Antibody (C-term) - Protein Information

Name TREX2

Function Exonuclease with a preference for double-stranded DNA with mismatched 3' termini. May play a role in DNA repair.

Cellular Location



Nucleus.

Tissue Location


Detected in heart, breast, prostate, skeletal muscle, testis, uterus, bone marrow, colon, small intestine, stomach and thymus.

TREX2 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

TREX2 Antibody (C-term) - Images



Western blot analysis of lysate from HL-60 cell line, using TREX2 Antibody (C-term)(Cat. #AP21082a). AP21082a was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysate at 20ug.

TREX2 Antibody (C-term) - Background

Exonuclease with a preference for double-stranded DNA with mismatched 3' termini. May play a role in DNA repair.

TREX2 Antibody (C-term) - References

Mazur D.J., et al.J. Biol. Chem. 274:19655-19660(1999). Mazur D.J., et al.J. Biol. Chem. 276:14718-14727(2001). Mazur D.J., et al.J. Biol. Chem. 276:17022-17029(2001). Perrino F.W., et al.J. Biol. Chem. 280:15212-15218(2005).