10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

(DANRE) adra2db Antibody (C-Term)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21170a

Specification

(DANRE) adra2db Antibody (C-Term) - Product Information

Application	**WB,E**
Primary Accession	Q8JG69
Reactivity	**Zebrafish**
Host	**Rabbit**
Clonality	**polyclonal**
Isotype	**Rabbit IgG**
Calculated MW	**46863**
Antigen Region	**323-356**

(DANRE) adra2db Antibody (C-Term) - Additional Information

Gene ID 266755

Other Names
Alpha-2Db adrenergic receptor, Alpha-2Db adrenoceptor, Alpha(2Db)AR, Alpha-2Db adrenoreceptor, adra2db

Target/Specificity
This DANRE adra2db antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 323-356 amino acids from the region of DANRE adra2db.

Dilution
WB~~1:4000

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
(DANRE) adra2db Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

(DANRE) adra2db Antibody (C-Term) - Protein Information

Name adra2db

Function Alpha-2 adrenergic receptors mediate the catecholamine- induced inhibition of adenylate cyclase through the action of G proteins. The order of potency for this receptor is

dexmedetomidine > norepinephrine = epinephrine > oxymetazoline.


Cellular Location
Cell membrane; Multi-pass membrane protein.

(DANRE) adra2db Antibody (C-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

(DANRE) adra2db Antibody (C-Term) - Images



All lanes : Anti-adra2db Antibody (C-Term) at 1:4000 dilution Lane 1: zebrafish muscle lysates Lane 2: Zebrafish lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 47 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

(DANRE) adra2db Antibody (C-Term) - Background

Alpha-2 adrenergic receptors mediate the catecholamine- induced inhibition of adenylate cyclase through the action of G proteins. The order of potency for this receptor is dexmedetomidine > norepinephrine = epinephrine > oxymetazoline.

(DANRE) adra2db Antibody (C-Term) - References

Ruuskanen J.O.,et al.Mol. Biol. Evol. 21:14-28(2004).
Ruuskanen J.O.,et al.Br. J. Pharmacol. 144:165-177(2005).