

UHRF1 Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21187a**Specification**

UHRF1 Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	Q96T88
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	89814

UHRF1 Antibody (N-term) - Additional Information**Gene ID** 29128**Other Names**

E3 ubiquitin-protein ligase UHRF1, 632-, Inverted CCAAT box-binding protein of 90 kDa, Nuclear protein 95, Nuclear zinc finger protein Np95, HuNp95, hNp95, RING finger protein 106, Transcription factor ICBP90, Ubiquitin-like PHD and RING finger domain-containing protein 1, hUHRF1, Ubiquitin-like-containing PHD and RING finger domains protein 1, UHRF1, ICBP90, NP95, RNF106

Target/Specificity

This UHRF1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 592-626 amino acids from the N-terminal region of human UHRF1.

Dilution

WB~~1:4000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

UHRF1 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

UHRF1 Antibody (N-term) - Protein Information**Name** UHRF1

Synonyms ICBP90, NP95, RNF106

Function Multidomain protein that acts as a key epigenetic regulator by bridging DNA methylation and chromatin modification. Specifically recognizes and binds hemimethylated DNA at replication forks via its YDG domain and recruits DNMT1 methyltransferase to ensure faithful propagation of the DNA methylation patterns through DNA replication. In addition to its role in maintenance of DNA methylation, also plays a key role in chromatin modification: through its tudor-like regions and PHD-type zinc fingers, specifically recognizes and binds histone H3 trimethylated at 'Lys-9' (H3K9me3) and unmethylated at 'Arg-2' (H3R2me0), respectively, and recruits chromatin proteins. Enriched in pericentric heterochromatin where it recruits different chromatin modifiers required for this chromatin replication. Also localizes to euchromatic regions where it negatively regulates transcription possibly by impacting DNA methylation and histone modifications. Has E3 ubiquitin-protein ligase activity by mediating the ubiquitination of target proteins such as histone H3 and PML. It is still unclear how E3 ubiquitin-protein ligase activity is related to its role in chromatin in vivo. Plays a role in DNA repair by cooperating with UHRF2 to ensure recruitment of FANCD2 to interstrand cross-links (ICLs) leading to FANCD2 activation. Acts as a critical player of proper spindle architecture by catalyzing the 'Lys-63'-linked ubiquitination of KIF11, thereby controlling KIF11 localization on the spindle (PubMed:[37728657](#)).

Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00358, ECO:0000269|PubMed:10646863, ECO:0000269|PubMed:17673620, ECO:0000269|PubMed:17967883, ECO:0000269|PubMed:19056828, ECO:0000269|PubMed:21777816, ECO:0000269|PubMed:30335751} Note=Associated, through the YDG domain (also called SRA domain), with replicating DNA from early to late S phase, including at replicating pericentric heterochromatin (By similarity). Also localizes to euchromatic regions. In non-S-phase cells, homogenously distributed through the nucleus (By similarity). {ECO:0000250|UniProtKB:Q8VDF2}


Tissue Location


Expressed in thymus, bone marrow, testis, lung and heart. Overexpressed in breast cancer.

UHRF1 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

UHRF1 Antibody (N-term) - Images



All lanes : Anti-UHRF1 Antibody (N-term) at 1:4000 dilution Lane 1: KG-1 whole cell lysates Lane 2: MCF-7 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 90 kDa Blocking/Dilution buffer: 5% NFDm/TBST.

UHRF1 Antibody (N-term) - Background

Multidomain protein that acts as a key epigenetic regulator by bridging DNA methylation and chromatin modification. Specifically recognizes and binds hemimethylated DNA at replication forks via its YDG domain and recruits DNMT1 methyltransferase to ensure faithful propagation of the DNA methylation patterns through DNA replication. In addition to its role in maintenance of DNA methylation, also plays a key role in chromatin modification: through its tudor-like regions and PHD-type zinc fingers, specifically recognizes and binds histone H3 trimethylated at 'Lys-9' (H3K9me3) and unmethylated at 'Arg-2' (H3R2me0), respectively, and recruits chromatin proteins. Enriched in pericentric heterochromatin where it recruits different chromatin modifiers required for this chromatin replication. Also localizes to euchromatic regions where it negatively regulates transcription possibly by impacting DNA methylation and histone modifications. Has E3 ubiquitin-protein ligase activity by mediating the ubiquitination of target proteins such as histone H3 and PML. It is still unclear how E3 ubiquitin-protein ligase activity is related to its role in chromatin in vivo. May be involved in DNA repair.

UHRF1 Antibody (N-term) - References

Hopfner R.,et al.Cancer Res. 60:121-128(2000).
Muto M.,et al.Radiat. Res. 166:723-733(2006).
Davenport J.W.,et al.Submitted (JUN-2000) to the EMBL/GenBank/DDBJ databases.
Bechtel S.,et al.BMC Genomics 8:399-399(2007).
Ota T.,et al.Nat. Genet. 36:40-45(2004).