10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

UBE2L6 Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP2118B

Specification

UBE2L6 Antibody (C-term) - Product Information

Application	**WB, IHC-P,E**
Primary Accession	O14933
Reactivity	**Human**
Host	**Rabbit**
Clonality	**Polyclonal**
Isotype	**Rabbit IgG**
Antigen Region	**123-153**

UBE2L6 Antibody (C-term) - Additional Information

Gene ID 9246

Other Names
Ubiquitin/ISG15-conjugating enzyme E2 L6, Retinoic acid-induced gene B protein, RIG-B, UbcH8, Ubiquitin carrier protein L6, Ubiquitin-protein ligase L6, UBE2L6, UBCH8

Target/Specificity
This UBE2L6 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 123-153 amino acids from the C-terminal region of human UBE2L6.

Dilution
WB~~1:1000
IHC-P~~1:10~50

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
UBE2L6 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

UBE2L6 Antibody (C-term) - Protein Information

Name UBE2L6

Synonyms UBCH8

Function Catalyzes the covalent attachment of ubiquitin or ISG15 to other proteins. Functions in the E6/E6-AP-induced ubiquitination of p53/TP53. Promotes ubiquitination and subsequent proteasomal degradation of FLT3.


Tissue Location
Present in natural killer cells (at protein level).

UBE2L6 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

UBE2L6 Antibody (C-term) - Images



Anti-UBE2L6 Antibody (K136) at 1:1000 dilution + MOLT-4 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 18 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999



Formalin-fixed and paraffin-embedded breast carcinoma tissue reacted with UBE2L6 Antibody (C-term)(Cat.#AP2118b), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

UBE2L6 Antibody (C-term) - Background

The modification of proteins with ubiquitin is an important cellular mechanism for targeting abnormal or short-lived proteins for degradation. Ubiquitination involves at least three classes of enzymes: ubiquitin-activating enzymes (E1s), ubiquitin-conjugating enzymes (E2s) and ubiquitin-protein ligases (E3s). UBE2L6 is a member of the E2 ubiquitin-conjugating enzyme family. This enzyme is demonstrated to function in the E6/E6-ap-induced ubiquitination of p53.

UBE2L6 Antibody (C-term) - References

Kumar, S., et al., J. Biol. Chem. 272(21):13548-13554 (1997). Ardley, H.C., et al., Cytogenet. Cell Genet. 89 (1-2), 137-140 (2000).

UBE2L6 Antibody (C-term) - Citations

- Influenza B virus non-structural protein 1 counteracts ISG15 antiviral activity by sequestering ISGylated viral proteins.
- Differential regulation of PML-RARÎ± stability by the ubiquitin ligases SIAH1/SIAH2 and TRIAD1.
- Mechanism for ubiquitylation of the leukemia fusion proteins AML1-ETO and PML-RARalpha.