

MAGI2 Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21243b

Specification

MAGI2 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	Q86UL8
Reactivity	Mouse, Rat
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	158754

MAGI2 Antibody (C-term) - Additional Information

Gene ID 9863

Other Names

Membrane-associated guanylate kinase, WW and PDZ domain-containing protein 2, Atrophin-1-interacting protein 1, AIP-1, Atrophin-1-interacting protein A, Membrane-associated guanylate kinase inverted 2, MAGI-2, MAGI2, ACVRINP1, AIP1, KIAA0705

Target/Specificity

This MAGI2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 1123-1156 amino acids from the C-terminal region of human MAGI2.

Dilution

WB~~1:2000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

MAGI2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

MAGI2 Antibody (C-term) - Protein Information

Name MAGI2

Synonyms ACVRINP1, AIP1, KIAA0705

Function Seems to act as a scaffold molecule at synaptic junctions by assembling neurotransmitter receptors and cell adhesion proteins (By similarity). Plays a role in nerve growth factor (NGF)-induced recruitment of RAPGEF2 to late endosomes and neurite outgrowth (By similarity). May play a role in regulating activin-mediated signaling in neuronal cells (By similarity). Enhances the ability of PTEN to suppress AKT1 activation (PubMed:[10760291](#)). Plays a role in receptor-mediated clathrin-dependent endocytosis which is required for ciliogenesis (By similarity).

Cellular Location

Cytoplasm. Late endosome. Synapse, synaptosome. Cell membrane; Peripheral membrane protein. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome {ECO:0000250|UniProtKB:Q9WVQ1}. Cell projection, cilium {ECO:0000250|UniProtKB:Q9WVQ1}. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome, centriole {ECO:0000250|UniProtKB:Q9WVQ1}. Photoreceptor inner segment {ECO:0000250|UniProtKB:Q9WVQ1}. Cell projection, cilium, photoreceptor outer segment {ECO:0000250|UniProtKB:Q9WVQ1}. Note=Localized diffusely in the cytoplasm before nerve growth factor (NGF) stimulation Recruited to late endosomes after NGF stimulation. Membrane-associated in synaptosomes (By similarity).

Tissue Location


Specifically expressed in brain.

MAGI2 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MAGI2 Antibody (C-term) - Images



All lanes : Anti-MAGI2 Antibody (C-term) at 1:2000 dilution Lane 1: mouse brain lysates Lane 2: rat brain lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 159 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

MAGI2 Antibody (C-term) - Background

Seems to act as scaffold molecule at synaptic junctions by assembling neurotransmitter receptors and cell adhesion proteins. May play a role in regulating activin-mediated signaling in neuronal cells. Enhances the ability of PTEN to suppress AKT1 activation. Plays a role in nerve growth factor (NGF)-induced recruitment of RAPGEF2 to late endosomes and neurite outgrowth.

MAGI2 Antibody (C-term) - References

Wood J.D.,et al.Mol. Cell. Neurosci. 11:149-160(1998).
Ishikawa K.,et al.DNA Res. 5:169-176(1998).
Hillier L.W.,et al.Nature 424:157-164(2003).
Scherer S.W.,et al.Science 300:767-772(2003).
Wu X.,et al.Proc. Natl. Acad. Sci. U.S.A. 97:4233-4238(2000).