10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

(DANRE) aplnra Antibody (C-Term)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21299b

Specification

(DANRE) aplnra Antibody (C-Term) - Product Information

Application	**WB,E**
Primary Accession	Q7SZP9
Reactivity	**Zebrafish**
Host	**Rabbit**
Clonality	**polyclonal**
Isotype	**Rabbit IgG**
Calculated MW	**40649**
Antigen Region	**343-377**

(DANRE) aplnra Antibody (C-Term) - Additional Information

Gene ID 561935

Other Names
Apelin receptor A, Angiotensin II receptor-like 1a, Angiotensin receptor-like 1a, G-protein coupled receptor APJ A, aplnra, agtrl1 {ECO:0000312|EMBL:ABI994701}, agtrl1a

Target/Specificity
This DANRE aplnra antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 343-377 amino acids from the human region of DANRE aplnra.

Dilution
WB~~1:2000

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
(DANRE) aplnra Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

(DANRE) aplnra Antibody (C-Term) - Protein Information

Name aplnra

Synonyms agtrl1 {ECO:0000312|EMBL:ABI99470.1}, ag

Function Receptor for apelin receptor early endogenous ligand (apela) and apelin (apln) hormones coupled to G proteins that inhibit adenylate cyclase activity (PubMed:17336906, PubMed:24316148, PubMed:24407481). Plays a key role in early development such as gastrulation, blood vessels formation and heart morphogenesis by acting as a receptor for apela hormone, promoting endoderm and mesendoderm cell migration and regulating the migration of cells fated to become myocardial progenitors, respectively (PubMed:17336906, PubMed:24316148, PubMed:24407481, PubMed:26017639). Positively regulates angioblast migration toward the embryonic midline, i.e. the position of the future vessel formation, during vasculogenesis (PubMed:26017639). May promote sinus venosus (SV)-derived endothelial cells migration into the developing heart to promote coronary blood vessel development (By similarity). Required for cardiovascular development, particularly for intersomitic vein angiogenesis by acting as a receptor for apln hormone (By similarity). Also plays a role in various processes in adults such as regulation of blood vessel formation, blood pressure, heart contractility, and heart failure (By similarity). Acts redundantly with agtrl1b in heart development (PubMed:17336906).

Cellular Location
Cell membrane {ECO:0000250|UniProtKB:P79960}; Multi-pass membrane protein {ECO:0000250|UniProtKB:P79960} Note=Internalized to the cytoplasm after exposure to apelin (apln) After exposure to apelin receptor early endogenous ligand (apela), internalized from the cell surface into an endosomal recycling compartment, from where it is recycled to the cell membrane {ECO:0000250|UniProtKB:P35414, ECO:0000250|UniProtKB:P79960}

Tissue Location
First expressed before epiboly in dorsal precursors. During epiboly, expressed in the enveloping layer, yolk syncytial layer and migrating mesendoderm. During segmentation stages, expressed in epithelial structures such as adaxial cells, border cells of the newly formed somites, developing lens, otic vesicles and venous vasculature.


(DANRE) aplnra Antibody (C-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

(DANRE) aplnra Antibody (C-Term) - Images

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999



All lanes : Anti-aplnra Antibody (C-Term) at 1:2000 dilution Lane 1: Zebrafish lysates Lane 2: Zebrafish muscle lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 41 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

(DANRE) aplnra Antibody (C-Term) - Background

Receptor for apelin coupled to G proteins that inhibit adenylate cyclase activity and plays a role in various processes in adults such as regulation of blood pressure, heart contractility, and heart failure. Also plays a key role in early development such as gastrulation and heart morphogenesis by acting as a receptor for apela hormone, promoting endoderm and mesendoderm cell migration and regulating the migration of cells fated to become myocardial progenitors, respectively (PubMed:24316148, PubMed:24407481). Acts redundantly with agtrl1b in heart development.

(DANRE) aplnra Antibody (C-Term) - References

Tucker B.,et al.Gene Expr. Patterns 7:258-265(2007).
Scott I.C.,et al.Dev. Cell 12:403-413(2007).
Chng S.C.,et al.Dev. Cell 27:672-680(2013).
Pauli A.,et al.Science 343:1248636-1248636(2014).