

(Mouse) Ctf1 Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21323a

Specification

(Mouse) Ctf1 Antibody - Product Information

Application	WB,E
Primary Accession	Q60753
Reactivity	Mouse, Rat
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	21509

(Mouse) Ctf1 Antibody - Additional Information

Gene ID 13019

Other Names

Cardiotrophin-1, CT-1, Ctf1

Target/Specificity

This Mouse Ctf1 antibody is generated from a rabbit immunized with a recombinant protein.

Dilution

WB~~1:2000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

(Mouse) Ctf1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

(Mouse) Ctf1 Antibody - Protein Information

Name Ctf1

Function Induces cardiac myocyte hypertrophy in vitro. Binds to and activates the ILST/gp130 receptor.

Cellular Location

Secreted.


Tissue Location

Highly expressed in heart, skeletal muscle, liver, lung and kidney. Lower levels in testis and brain.
No expression in spleen


(Mouse) Ctf1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

(Mouse) Ctf1 Antibody - Images

Anti-Ctf1 Antibody at 1:2000 dilution + rat lung lysates. Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 22 kDa. Blocking/Dilution buffer: 5% NFDM/TBST.



All lanes : Anti-Ctf1 Antibody at 1:2000 dilution + mouse skeletal muscle lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 22 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

(Mouse) Ctf1 Antibody - Background

Induces cardiac myocyte hypertrophy in vitro. Binds to and activates the ILST/gp130 receptor.

(Mouse) Ctf1 Antibody - References

Pennica D.,et al.Proc. Natl. Acad. Sci. U.S.A. 92:1142-1146(1995).