

**NBPF12 Antibody (Center)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP21456c****Specification**

---

**NBPF12 Antibody (Center) - Product Information**

Application	WB,E
Primary Accession	<a href="#">Q5TAG4</a>
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	167689
Antigen Region	148-181

**NBPF12 Antibody (Center) - Additional Information****Gene ID** 149013**Other Names**

Neuroblastoma breakpoint family member 12, Chromosome 1 amplified sequence 1, NBPF12, COAS1, KIAA1245

**Target/Specificity**

This NBPF12 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 148-181 amino acids from the Central region of human NBPF12.

**Dilution**

WB~~1:2000

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

NBPF12 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

**NBPF12 Antibody (Center) - Protein Information****Name** NBPF12**Synonyms** COAS1, KIAA1245

**Cellular Location**

Cytoplasm.


**Tissue Location**

Widely expressed with highest levels in brain, ovary, mammary gland, skin and adipose tissue. Also expressed in testis. Detected in a number of tumors including osteosarcoma, mammary carcinoma and hepatocellular carcinoma.

**NBPF12 Antibody (Center) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**NBPF12 Antibody (Center) - Images**

Anti-NBPF12 Antibody (Center) at 1:2000 dilution + human ovary lysates. Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 31 kDa. Blocking/Dilution buffer: 5% NFDM/TBST.

**NBPF12 Antibody (Center) - References**

- Gregory S.G., et al. Nature 441:315-321(2006).  
Mei G., et al. Submitted (FEB-1999) to the EMBL/GenBank/DDBJ databases.  
Meza-Zepeda L.A., et al. Oncogene 21:2261-2269(2002).  
Vandepoele K., et al. Mol. Biol. Evol. 22:2265-2274(2005).