

C17orf80 Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21485c**Specification**

C17orf80 Antibody (Center) - Product Information

Application	WB,E
Primary Accession	Q9BSJ5
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	67315
Antigen Region	264-696

C17orf80 Antibody (Center) - Additional Information**Gene ID** 55028**Other Names**

Uncharacterized protein C17orf80, Cell migration-inducing gene 3 protein, Human lung cancer oncogene 8 protein, HLC-8, C17orf80

Target/Specificity

This C17orf80 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 264-696 amino acids from the Central region of human C17orf80.

Dilution

WB~~1:1000-1:2000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

C17orf80 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

C17orf80 Antibody (Center) - Protein Information**Name** MTNAP1 ([HGNC:29601](#))

Function Critical regulator of mitochondrial DNA (mtDNA) abundance (PubMed:[37676315](#)). Binds dsDNA throughout the mitochondrial genome without sequence specificity and controls mtDNA

copy number by promoting its replication (PubMed:[37676315](#)). Also plays important roles in mitochondrial metabolism and cell proliferation (PubMed:[37676315](#)).

Cellular Location


Mitochondrion inner membrane; Single-pass type I membrane protein. Mitochondrion matrix, mitochondrion nucleoid Note=Retained in nucleoids under mtDNA replication stress caused by ddC and EtBr.

C17orf80 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

C17orf80 Antibody (Center) - Images



All lanes : Anti-C17orf80 Antibody (Center) at 1:1000-1:2000 dilution Lane 1: K562 whole cell lysates Lane 2: human ovary lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 67 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

C17orf80 Antibody (Center) - References

Kim J.W.,et al.Submitted (OCT-2002) to the EMBL/GenBank/DDBJ databases.
Wiemann S.,et al.Genome Res. 11:422-435(2001).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Zody M.C.,et al.Nature 440:1045-1049(2006).
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.