

**HNRNPUL2 Antibody (Center)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP21566c****Specification**

---

**HNRNPUL2 Antibody (Center) - Product Information**

Application	WB, IHC-P,E
Primary Accession	<a href="#">Q1KMD3</a>
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	85105

**HNRNPUL2 Antibody (Center) - Additional Information****Gene ID** 221092**Other Names**

Heterogeneous nuclear ribonucleoprotein U-like protein 2, Scaffold-attachment factor A2, SAF-A2, HNRNPUL2, HNRPUL2

**Target/Specificity**

This HNRNPUL2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 337-350 amino acids from the Central region of human HNRNPUL2.

**Dilution**

WB~~1:2000

IHC-P~~1:25

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

HNRNPUL2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

**HNRNPUL2 Antibody (Center) - Protein Information****Name** HNRNPUL2**Synonyms** HNRPUL2


**Cellular Location**

Nucleus {ECO:0000250|UniProtKB:Q00PI9}.


**HNRNPUL2 Antibody (Center) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**HNRNPUL2 Antibody (Center) - Images**

Anti-HNRNPUL2 Antibody (Center) at 1:8000 dilution + Hela whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 85 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



All lanes : Anti-HNRNPUL2 Antibody (Center) at 1:2000 dilution Lane 1: MCF-7 whole cell lysates Lane 2: Jurkat whole cell lysates Lane 3: 293T/17 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 85 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



AP21566c staining HNRNPUL2 in human stomach tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hour at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.

#### **HNRNPUL2 Antibody (Center) - References**

Herrmann F.,et al.Submitted (MAR-2006) to the EMBL/GenBank/DDBJ databases.  
Bechtel S.,et al.BMC Genomics 8:399-399(2007).  
Bienvenut W.V.,et al.Submitted (DEC-2008) to UniProtKB.  
Burkard T.R.,et al.BMC Syst. Biol. 5:17-17(2011).