10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

EPS15 Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP2159b

Specification

EPS15 Antibody (C-term) - Product Information

Application	**WB,E**
Primary Accession	P42566
Other Accession	P42567, NP_001972
Reactivity	**Human**
Predicted	**Mouse**
Host	**Rabbit**
Clonality	**Polyclonal**
Isotype	**Rabbit IgG**
Calculated MW	**98656**
Antigen Region	**840-870**

EPS15 Antibody (C-term) - Additional Information

Gene ID 2060

Other Names
Epidermal growth factor receptor substrate 15, Protein Eps15, Protein AF-1p, EPS15, AF1P

Target/Specificity
This EPS15 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 840-870 amino acids from the C-terminal region of human EPS15.

Dilution
WB~~1:1000

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
EPS15 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

EPS15 Antibody (C-term) - Protein Information

Name EPS15

Synonyms AF1P

Function Involved in cell growth regulation. May be involved in the regulation of mitogenic signals and control of cell proliferation. Involved in the internalization of ligand-inducible receptors of the receptor tyrosine kinase (RTK) type, in particular EGFR. Plays a role in the assembly of clathrin-coated pits (CCPs). Acts as a clathrin adapter required for post-Golgi trafficking. Seems to be involved in CCPs maturation including invagination or budding. Involved in endocytosis of integrin beta-1 (ITGB1) and transferrin receptor (TFR); internalization of ITGB1 as DAB2-dependent cargo but not TFR seems to require association with DAB2.

Cellular Location
Cytoplasm. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Membrane, clathrin-coated pit Note=Recruited to the plasma membrane upon EGFR activation and localizes to coated pits. Colocalizes with UBQLN1 in ubiquitin-rich cytoplasmic aggregates that are not endocytic compartments and in cytoplasmic juxtanuclear structures called aggresomes


Tissue Location
Ubiquitously expressed.

EPS15 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

EPS15 Antibody (C-term) - Images



Western blot analysis of anti-EPS15 Pab (Cat. #AP2159b) in HL60 cell line lysate (35ug/lane). EPS15 (arrow) was detected using the purified Pab.

EPS15 Antibody (C-term) - Background

EPS15 is involved in cell growth regulation, possibly via the regulation of mitogenic signals and control of cell proliferation. EPS15 also participates in the internalization of ligand-inducible receptors of the receptor tyrosine kinase (RTK) type, in particular EGFR. Potential interaction partners include AP2A2, STN2, EPN1, and CRK via its SH3-binding sites. EPS15 is ubiquitously expressed. Phosphorylation on Tyr-849 is involved in the internalization of EGFR. EPS15 is not required for membrane translocation after EGF treatment or for targeting to coated pits, but essential for a subsequent step in EGFR endocytosis. This protein is involved in a t(1;11)(p32;q23) chromosomal translocation in acute leukemias causing fusion to the trithorax (MLL or HRX) gene product which contains DNA-binding motifs resulting in a rogue activator protein. Structurally, EPS15 contains 2 EF-hand calcium-binding domains, 3 EH domains, and 2 ubiquitin-interacting motif (UIM) repeats.

EPS15 Antibody (C-term) - References

de Beer, T., et al., Nat. Struct. Biol. 7(11):1018-1022 (2000).
Enmon, J.L., et al., Biochemistry 39(15):4309-4319 (2000).
de Beer, T., et al., Science 281(5381):1357-1360 (1998).
Chen, H., et al., Nature 394(6695):793-797 (1998).
Matsuda, M., et al., J. Biol. Chem. 271(24):14468-14472 (1996).