

Mouse Srms Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP21603c

Specification

Mouse Srms Antibody (Center) - Product Information

Application WB,E **Primary Accession** Q62270 Reactivity Mouse Host **Rabbit** Clonality polyclonal Isotype Rabbit IgG Calculated MW 55758 Antigen Region 113-147

Mouse Srms Antibody (Center) - Additional Information

Other Names

Tyrosine-protein kinase Srms, PTK70, Srms, Srm

Target/Specificity

This Mouse Srms antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 113-147 amino acids from the Central region of Mouse Srms.

Dilution

WB~~1:2000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Mouse Srms Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Mouse Srms Antibody (Center) - Protein Information

Name Srms

Synonyms Srm

Function Non-receptor tyrosine-protein kinase which phosphorylates DOK1 on tyrosine residues. Also phosphorylates KHDRBS1/SAM68 and VIM on tyrosine residues. Phosphorylation of KHDRBS1 is EGF-dependent. Phosphorylates OTUB1, promoting deubiquitination of RPTOR.



Cellular Location

Cytoplasm {ECO:0000250|UniProtKB:Q9H3Y6}. Note=Localizes to punctate cytoplasmic structures {ECO:0000250|UniProtKB:Q9H3Y6}

Tissue Location


Higher expression in liver, lung, thymus and skin than in brain, kidney, heart and spleen (PubMed:9226137). In skin, highly expressed in keratinocytes (PubMed:9226137). Abundant in lung, liver, spleen, kidney and testis and is also detected in the cerebrum (PubMed:7935409).

Mouse Srms Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Mouse Srms Antibody (Center) - Images



Anti-Srms Antibody (Center)at 1:2000 dilution + mouse spleen lysates Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 56 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Mouse Srms Antibody (Center) - Background

Non-receptor tyrosine-protein kinase which phosphorylates DOK1 on tyrosine residues. May be involved in proliferation or differentiation of keratinocytes in the skin.

Mouse Srms Antibody (Center) - References

Kawachi Y., et al. Exp. Dermatol. 6:140-146(1997). Kohmura N., et al. Mol. Cell. Biol. 14:6915-6925(1994).





Wall M., et al. Submitted (APR-2001) to the EMBL/GenBank/DDBJ databases.