10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

WIPF3 Antibody (N-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21633a

Specification

WIPF3 Antibody (N-term) - Product Information

Application	**WB,E**
Primary Accession	A6NGB9
Reactivity	**Human, Rat**
Host	**Rabbit**
Clonality	**polyclonal**
Isotype	**Rabbit IgG**
Calculated MW	**49458**

WIPF3 Antibody (N-term) - Additional Information

Gene ID 644150

Other Names
WAS/WASL-interacting protein family member 3, Corticosteroids and regional expression protein 16 homolog, WIPF3, CR16

Target/Specificity
This WIPF3 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 109-142 amino acids from the N-terminal region of human WIPF3.

Dilution
WB~~1:2000

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
WIPF3 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

WIPF3 Antibody (N-term) - Protein Information

Name WIPF3

Synonyms CR16

Function May be a regulator of cytoskeletal organization. May have a role in spermatogenesis (By

similarity).


Cellular Location
Cytoplasm. Note=In hippocampal neurons colocalizes with WASL in the cell body, axons and the growth cone

WIPF3 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

WIPF3 Antibody (N-term) - Images



All lanes : Anti-WIPF3 Antibody (N-term) at 1:2000 dilution Lane 1: human liver lysate Lane 2: rat liver lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 49 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

WIPF3 Antibody (N-term) - Background

May be a regulator of cytoskeletal organization. May have a role in spermatogenesis (By similarity).

WIPF3 Antibody (N-term) - References

Hillier L.W.,et al.Nature 424:157-164(2003).
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.
Ebert L.,et al.Submitted (JAN-2003) to the EMBL/GenBank/DDBJ databases.
Zettl M.,et al.Curr. Biol. 12:1617-1622(2002).

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999