

TST Antibody (C-Term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21669b**Specification**

TST Antibody (C-Term) - Product Information

Application	WB,E
Primary Accession	Q16762
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	33429

TST Antibody (C-Term) - Additional Information**Gene ID** 7263**Other Names**

Thiosulfate sulfurtransferase, Rhodanese, TST

Target/Specificity

This TST antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 208-232 amino acids from the human TST.

Dilution

WB~~1:2000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

TST Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

TST Antibody (C-Term) - Protein Information**Name** TST

Function Formation of iron-sulfur complexes, cyanide detoxification or modification of sulfur-containing enzymes. Other thiol compounds, besides cyanide, can act as sulfur ion acceptors. Also has weak mercaptopyruvate sulfurtransferase (MST) activity (By similarity). Together with MRPL18, acts as a mitochondrial import factor for the cytosolic 5S rRNA. Only the

nascent unfolded cytoplasmic form is able to bind to the 5S rRNA.

Cellular Location


Mitochondrion matrix.

TST Antibody (C-Term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TST Antibody (C-Term) - Images



All lanes : Anti-TST Antibody (C-Term) at 1:8000 dilution Lane 1: human liver lysate Lane 2: HepG2 lysate Lane 3: mouse liver lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 33 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Anti-TST Antibody (C-Term) at 1:2000 dilution + rat kidney lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 33 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

TST Antibody (C-Term) - Background

Formation of iron-sulfur complexes, cyanide detoxification or modification of sulfur-containing enzymes. Other thiol compounds, besides cyanide, can act as sulfur ion acceptors. Also has weak mercaptopyruvate sulfurtransferase (MST) activity (By similarity). Together with MRPL18, acts as a mitochondrial import factor for the cytosolic 5S rRNA. Only the nascent unfolded cytoplasmic form is able to bind to the 5S rRNA.

TST Antibody (C-Term) - References

Aita N.,et al.Biochem. Biophys. Res. Commun. 231:56-60(1997).
Collins J.E.,et al.Genome Biol. 5:R84.1-R84.11(2004).
Ebert L.,et al.Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Dunham I.,et al.Nature 402:489-495(1999).