

GMCL1 Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21774c**Specification**

GMCL1 Antibody (Center) - Product Information

Application	WB, IHC-P,E
Primary Accession	Q96IK5
Reactivity	Human, Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	58685

GMCL1 Antibody (Center) - Additional Information**Gene ID** 64395**Other Names**

Germ cell-less protein-like 1, GMCL1, GCL

Target/Specificity

This GMCL1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 281-315 amino acids from the Central region of human GMCL1.

Dilution

WB~~1:2000

IHC-P~~1:25

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

GMCL1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

GMCL1 Antibody (Center) - Protein Information**Name** GMCL1**Synonyms** BTBD13, GCL, SPATA29**Function** Possible function in spermatogenesis. Enhances the degradation of MDM2 and increases

the amount of p53 probably by modulating the nucleocytoplasmic transport (By similarity).

Cellular Location


Nucleus matrix.

GMCL1 Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

GMCL1 Antibody (Center) - Images



All lanes : Anti-GMCL1 Antibody (Center) at 1:2000 dilution Lane 1: 293T/17 whole cell lysate Lane 2: HL-60 whole cell lysate Lane 3: mouse testis lysate Lane 4: MCF-7 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 59 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



AP21774c staining GMCL1 in human kidney tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hour at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.

GMCL1 Antibody (Center) - Background

Possible function in spermatogenesis. Enhances the degradation of MDM2 and increases the amount of p53 probably by modulating the nucleocytoplasmic transport (By similarity).

GMCL1 Antibody (Center) - References

Ota T., et al. Nat. Genet. 36:40-45(2004).
Holaska J.M., et al. J. Biol. Chem. 278:6969-6975(2003).