

PLA2G4D Antibody (C-Term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21807b

Specification

PLA2G4D Antibody (C-Term) - Product Information

Application	WB,E
Primary Accession	Q86XP0
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	91952
Antigen Region	754-786

PLA2G4D Antibody (C-Term) - Additional Information

Gene ID 283748

Other Names

Cytosolic phospholipase A2 delta, cPLA2-delta, Phospholipase A2 group IVD, PLA2G4D

Target/Specificity

This PLA2G4D antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 754-786 amino acids from human PLA2G4D.

Dilution

WB~~1:2000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PLA2G4D Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

PLA2G4D Antibody (C-Term) - Protein Information

Name PLA2G4D ([HGNC:30038](#))

Function Calcium-dependent phospholipase A2 that selectively hydrolyzes glycerophospholipids in the sn-2 position (PubMed:[14709560](#)). Has a preference for linoleic acid at the sn-2 position (PubMed:[14709560](#)).

Cellular Location

Cytoplasm, cytosol {ECO:0000250|UniProtKB:Q50L43}. Membrane

{ECO:0000250|UniProtKB:Q50L43}; Peripheral membrane protein

{ECO:0000250|UniProtKB:Q50L43}; Cytoplasmic side {ECO:0000250|UniProtKB:Q50L43}.

Note=Translocates to perinuclear membranes that may correspond to endoplasmic reticulum or Golgi in a calcium-dependent fashion. {ECO:0000250|UniProtKB:Q50L43}

Tissue Location

Expressed in stratified squamous epithelia, such as those in skin and cervix, but not in other

tissues (PubMed:14709560) Strongly expressed in the upper spinous layer of the psoriatic


epidermis, expressed weakly and discontinuously in atopic dermatitis and mycosis fungoides, and

not detected in the epidermis of normal skin (PubMed:14709560).

PLA2G4D Antibody (C-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PLA2G4D Antibody (C-Term) - Images

All lanes : Anti-PLA2G4D Antibody (C-Term) at 1:2000 dilution Lane 1: A431 whole cell lysate Lane 2: A549 whole cell lysate Lane 3: HeLa whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 92 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

PLA2G4D Antibody (C-Term) - Background

Calcium-dependent phospholipase A2 that selectively hydrolyzes glycerophospholipids in the sn-2 position. Not arachidonic acid-specific but has linoleic acid-specific activity. May play a role in

inflammation in psoriatic lesions.

PLA2G4D Antibody (C-Term) - References

Chiba H.,et al.J. Biol. Chem. 279:12890-12897(2004).

Zody M.C.,et al.Nature 440:671-675(2006).

Tao R.,et al.Prostaglandins Leukot. Essent. Fatty Acids 73:419-422(2005).