

## **BBC3 Antibody (C-Term)**

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21821b

## **Specification**

#### **BBC3 Antibody (C-Term) - Product Information**

WB,E Application **Primary Accession 096PG8** Reactivity Human Host Rabbit Clonality polyclonal Isotype Rabbit IgG Calculated MW 26498 Antigen Region 158-192

## **BBC3 Antibody (C-Term) - Additional Information**

#### **Gene ID 27113**

#### **Other Names**

Bcl-2-binding component 3, JFY-1, p53 up-regulated modulator of apoptosis, BBC3, PUMA

# **Target/Specificity**

This BBC3 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 158-192 amino acids from human BBC3.

#### **Dilution**

WB~~1:2000

#### **Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

## **Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### **Precautions**

BBC3 Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

## **BBC3 Antibody (C-Term) - Protein Information**

#### Name BBC3

## **Synonyms PUMA**

Function [Isoform 3]: Does not affect cell growth.



**Cellular Location** 


Note=Contrary to isoforms 1 and 2, isoform 3 does not localize to the mitochondria

## **BBC3 Antibody (C-Term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

# **BBC3 Antibody (C-Term) - Images**



Anti-BBC3 Antibody (C-Term) at 1:2000 dilution + A549 whole cell lysate Lysates/proteins at 20  $\mu$ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 26 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

# BBC3 Antibody (C-Term) - Background

Essential mediator of p53/TP53-dependent and p53/TP53- independent apoptosis. Isoform 3 fails to show any growth- inhibitory or apoptotic activity.

# **BBC3 Antibody (C-Term) - References**

Nakano K., et al. Mol. Cell 7:683-694(2001). Grimwood J., et al. Nature 428:529-535(2004). Mural R.J., et al. Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.