10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

ZBTB7C Antibody (N-Term)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21825a

Specification

ZBTB7C Antibody (N-Term) - Product Information

Application	**WB,E**
Primary Accession	A1YPR0
Reactivity	**Human**
Host	**Rabbit**
Clonality	**polyclonal**
Isotype	**Rabbit IgG**
Calculated MW	**69017**
Antigen Region	**168-198**

ZBTB7C Antibody (N-Term) - Additional Information

Gene ID 201501

Other Names
Zinc finger and BTB domain-containing protein 7C, Affected by papillomavirus DNA integration in ME180 cells protein 1, APM-1, Zinc finger and BTB domain-containing protein 36, Zinc finger protein 857C, ZBTB7C, APM1, ZBTB36, ZNF857C

Target/Specificity
This ZBTB7C antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 168-198 amino acids from human ZBTB7C.

Dilution
WB~~1:2000

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
ZBTB7C Antibody (N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

ZBTB7C Antibody (N-Term) - Protein Information

Name ZBTB7C

Synonyms APM1, ZBTB36, ZNF857C

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Function May be a tumor suppressor gene.


Tissue Location
Detected in normal cervical keratinocytes, and in some cervical carcinoma cell lines.

ZBTB7C Antibody (N-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

ZBTB7C Antibody (N-Term) - Images



All lanes : Anti-ZBTB7C Antibody (N-Term) at 1:2000 dilution Lane 1: human placenta lysate Lane 2: human uterus lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 69 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

ZBTB7C Antibody (N-Term) - Background

May be a tumor suppressor gene.

ZBTB7C Antibody (N-Term) - References

Reuter S.,et al.EMBO J. 17:215-222(1998).
Gilling M.,et al.Submitted (NOV-2006) to the EMBL/GenBank/DDBJ databases.