

NQO2 Antibody (N-Term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21839a**Specification**

NQO2 Antibody (N-Term) - Product Information

Application	WB,E
Primary Accession	P16083
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	25919

NQO2 Antibody (N-Term) - Additional Information**Gene ID** 4835**Other Names**

Ribosyldihydronicotinamide dehydrogenase [quinone], NRH dehydrogenase [quinone] 2, NRH:quinone oxidoreductase 2, Quinone reductase 2, QR2, NQO2, NMOR2

Target/Specificity

This NQO2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 21-52 amino acids from human NQO2.

Dilution

WB~~1:2000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

NQO2 Antibody (N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

NQO2 Antibody (N-Term) - Protein Information**Name** NQO2**Synonyms** NMOR2**Function** The enzyme apparently serves as a quinone reductase in connection with conjugation

reactions of hydroquinones involved in detoxification pathways as well as in biosynthetic processes such as the vitamin K-dependent gamma-carboxylation of glutamate residues in prothrombin synthesis.

Cellular Location


Cytoplasm.

NQO2 Antibody (N-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NQO2 Antibody (N-Term) - Images



All lanes : Anti-NQO2 Antibody (N-Term) at 1:2000 dilution Lane 1: 293T/17 whole cell lysate Lane 2: HeLa whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 26 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

NQO2 Antibody (N-Term) - Background

The enzyme apparently serves as a quinone reductase in connection with conjugation reactions of hydroquinones involved in detoxification pathways as well as in biosynthetic processes such as the vitamin K-dependent gamma-carboxylation of glutamate residues in prothrombin synthesis.

NQO2 Antibody (N-Term) - References

Jaiswal A.K.,et al.Biochemistry 29:1899-1906(1990).
Jaiswal A.K.,et al.J. Biol. Chem. 269:14502-14508(1994).

Iida A.,et al.Submitted (OCT-2000) to the EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Mungall A.J.,et al.Nature 425:805-811(2003).