

MYBL2 Antibody (C-Term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21853b**Specification**

MYBL2 Antibody (C-Term) - Product Information

Application	WB,E
Primary Accession	P10244
Reactivity	Human, Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	78764

MYBL2 Antibody (C-Term) - Additional Information**Gene ID** 4605**Other Names**

Myb-related protein B, B-Myb, Myb-like protein 2, MYBL2, BMYB

Target/Specificity

This MYBL2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 540-571 amino acids from human MYBL2.

Dilution

WB~~1:8000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

MYBL2 Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

MYBL2 Antibody (C-Term) - Protein Information**Name** MYBL2**Synonyms** BMYB

Function Transcription factor involved in the regulation of cell survival, proliferation, and differentiation. Transactivates the expression of the CLU gene.


Cellular Location

Nucleus.


MYBL2 Antibody (C-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MYBL2 Antibody (C-Term) - Images

All lanes : Anti-MYBL2 Antibody (C-Term) at 1:2000 dilution Lane 1: 293T/17 whole cell lysate Lane 2: K562 whole cell lysate Lane 3: HeLa whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 79 kDa Blocking/Dilution buffer: 5% NFDm/TBST.



All lanes : Anti-MYBL2 Antibody (C-Term) at 1:8000 dilution Lane 1: CEM whole cell lysate Lane 2: MCF-7 whole cell lysate Lane 3: mouse testis lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 79 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

MYBL2 Antibody (C-Term) - Background

Transcription factor involved in the regulation of cell survival, proliferation, and differentiation. Transactivates the expression of the CLU gene.

MYBL2 Antibody (C-Term) - References

Nomura N.,et al.Nucleic Acids Res. 16:11075-11089(1988).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Totoki Y.,et al.Submitted (APR-2005) to the EMBL/GenBank/DDBJ databases.
Deloukas P.,et al.Nature 414:865-871(2001).
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.