10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

C18orf32 Antibody (C-Term)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21952b

Specification

C18orf32 Antibody (C-Term) - Product Information

Application	**WB,E**
Primary Accession	Q8TCD1
Reactivity	**Human**
Host	**Rabbit**
Clonality	**polyclonal**
Isotype	**Rabbit IgG**
Calculated MW	**8669**

C18orf32 Antibody (C-Term) - Additional Information

Gene ID 497661

Other Names
UPF0729 protein C18orf32, Putative NF-kappa-B-activating protein 200, C18orf32

Target/Specificity
This C18orf32 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 26-58 amino acids from human C18orf32.

Dilution
WB~~1:2000

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
C18orf32 Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

C18orf32 Antibody (C-Term) - Protein Information

Name C18orf32

Function May activate the NF-kappa-B signaling pathway.

Cellular Location
Endoplasmic reticulum. Lipid droplet


10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

C18orf32 Antibody (C-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

C18orf32 Antibody (C-Term) - Images



All lanes : Anti-C18orf32 Antibody (C-Term) at 1:2000 dilution Lane 1: human brain lysate Lane 2: LNCaP whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 9 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

C18orf32 Antibody (C-Term) - Background

May activate the NF-kappa-B signaling pathway.

C18orf32 Antibody (C-Term) - References

Matsuda A.,et al.Oncogene 22:3307-3318(2003).
Ota T.,et al.Nat. Genet. 36:40-45(2004).