

## TAX1BP3 Antibody (C-Term)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP22005b

## **Specification**

### TAX1BP3 Antibody (C-Term) - Product Information

Application WB,E
Primary Accession 014907

Other Accession <u>Q9DBG9</u>, <u>Q4QQV1</u>

Reactivity
Predicted
Host
Clonality
Isotype
Calculated MW
Human
Mouse, Rat
Rabbit
Rabbit
Rabbit
Polyclonal
Rabbit IgG
13735

## TAX1BP3 Antibody (C-Term) - Additional Information

#### Gene ID 30851

#### **Other Names**

Tax1-binding protein 3, Glutaminase-interacting protein 3, Tax interaction protein 1, TIP-1, Tax-interacting protein 1, TAX1BP3 (<a href="http://www.genenames.org/cgi-bin/gene\_symbol\_report?hgnc\_id=30684" target="blank">HGNC:30684</a>)

## Target/Specificity

This TAX1BP3 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 71-103 amino acids from human TAX1BP3.

# **Dilution**

WB~~1:2000

## **Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### **Precautions**

TAX1BP3 Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

## TAX1BP3 Antibody (C-Term) - Protein Information

## Name TAX1BP3 (HGNC:30684)

**Function** May regulate a number of protein-protein interactions by competing for PDZ domain binding sites. Binds CTNNB1 and may thereby act as an inhibitor of the Wnt signaling pathway. Competes with LIN7A for KCNJ4 binding, and thereby promotes KCNJ4 internalization. May play a



role in the Rho signaling pathway. May play a role in activation of CDC42 by the viral protein HPV16 E6.

#### **Cellular Location**

Cytoplasm. Nucleus. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Note=Recruited to the cell membrane by interaction with membrane proteins

#### **Tissue Location**


Ubiquitous. Detected in brain, heart, kidney, lung, small intestine and skeletal muscle. Detected in various cell lines including HeLa. Weakly expressed in peripheral blood leukocytes

## TAX1BP3 Antibody (C-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

## TAX1BP3 Antibody (C-Term) - Images



Anti-TAX1BP3 Antibody (C-Term) at 1:2000 dilution + HT-1080 whole cell lysate Lysates/proteins at 20  $\mu$ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 14 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

## TAX1BP3 Antibody (C-Term) - Background

May regulate a number of protein-protein interactions by competing for PDZ domain binding sites. Binds CTNNB1 and may thereby act as an inhibitor of the Wnt signaling pathway. Competes with LIN7A for KCNJ4 binding, and thereby promotes KCNJ4 internalization. May play a role in the Rho signaling pathway. May play a role in activation of CDC42 by the viral protein HPV16 E6.



# TAX1BP3 Antibody (C-Term) - References

Rousset R.,et al.Oncogene 16:643-654(1998). Touchman J.W.,et al.Genome Res. 10:165-173(2000). Olalla L.,et al.FEBS Lett. 488:116-122(2001). Honore B.,et al.Submitted (JUN-2000) to the EMBL/GenBank/DDBJ databases. Ota T.,et al.Nat. Genet. 36:40-45(2004).