

COPS7B Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP22164c**Specification**

COPS7B Antibody (Center) - Product Information

Application	WB,E
Primary Accession	Q9H9Q2
Other Accession	Q2KI56
Reactivity	Human
Predicted	Bovine
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	29622

COPS7B Antibody (Center) - Additional Information**Gene ID** 64708**Other Names**

COP9 signalosome complex subunit 7b, SGN7b, Signalosome subunit 7b, JAB1-containing signalosome subunit 7b, COPS7B, CSN7B

Target/Specificity

This COPS7B antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 223-257 amino acids from the Central region of human COPS7B.

Dilution

WB~~1:500-1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

COPS7B Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

COPS7B Antibody (Center) - Protein Information**Name** COPS7B**Synonyms** CSN7B

Function Component of the COP9 signalosome complex (CSN), a complex involved in various cellular and developmental processes. The CSN complex is an essential regulator of the ubiquitin (Ubl) conjugation pathway by mediating the deneddylation of the cullin subunits of SCF-type E3 ligase complexes, leading to decrease the Ubl ligase activity of SCF-type complexes such as SCF, CSA or DDB2. The complex is also involved in phosphorylation of p53/TP53, JUN, I-kappa-B-alpha/NFKBIA, ITPK1 and IRF8/ICSBP, possibly via its association with CK2 and PKD kinases. CSN-dependent phosphorylation of TP53 and JUN promotes and protects degradation by the Ubl system, respectively.

Cellular Location


Cytoplasm. Nucleus.

COPS7B Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

COPS7B Antibody (Center) - Images



All lanes : Anti-COPS7B Antibody (Center) at 1:500-1:1000 dilution Lane 1: Jurkat whole cell lysate Lane 2: HL-60 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 30 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

COPS7B Antibody (Center) - Background

Component of the COP9 signalosome complex (CSN), a complex involved in various cellular and developmental processes. The CSN complex is an essential regulator of the ubiquitin (Ubl) conjugation pathway by mediating the deneddylation of the cullin subunits of SCF-type E3 ligase

complexes, leading to decrease the Ubl ligase activity of SCF-type complexes such as SCF, CSA or DDB2. The complex is also involved in phosphorylation of p53/TP53, JUN, I-kappa-B-alpha/NFKBIA, ITPK1 and IRF8/ICSBP, possibly via its association with CK2 and PKD kinases. CSN-dependent phosphorylation of TP53 and JUN promotes and protects degradation by the Ubl system, respectively.

COPS7B Antibody (Center) - References

Ota T.,et al.Nat. Genet. 36:40-45(2004).
Hillier L.W.,et al.Nature 434:724-731(2005).
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.
Bech-Otschir D.,et al.EMBO J. 20:1630-1639(2001).
Lyapina S.,et al.Science 292:1382-1385(2001).