

**FAM159B Antibody (C-Term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP22187b**

**Specification**

---

**FAM159B Antibody (C-Term) - Product Information**

|                   |                        |
|-------------------|------------------------|
| Application       | WB,E                   |
| Primary Accession | <a href="#">A6NKG6</a> |
| Reactivity        | Human                  |
| Host              | Rabbit                 |
| Clonality         | polyclonal             |
| Isotype           | Rabbit IgG             |
| Antigen Region    | 121-154                |

**FAM159B Antibody (C-Term) - Additional Information**

**Gene ID** 100132916

**Target/Specificity**

This FAM159B antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 121-154 amino acids from human FAM159B.

**Dilution**

WB~~1:2000

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

FAM159B Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

**FAM159B Antibody (C-Term) - Protein Information**

**Name** SHISAL2B ([HGNC:34236](#))

**Synonyms** FAM159B

**Cellular Location**


Membrane; Single-pass membrane protein

## FAM159B Antibody (C-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## FAM159B Antibody (C-Term) - Images



All lanes : Anti-FAM159B Antibody (C-Term) at 1:2000 dilution Lane 1: MKN45 whole cell lysate Lane 2: PANC-1 whole cell lysate Lysates/proteins at 20  $\mu$ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 18 kDa Blocking/Dilution buffer: 5% NFDM/TBST.