10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

NSFL1C Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP22236c

Specification

NSFL1C Antibody (Center) - Product Information

Application	**WB, IHC-P, FC,E**
Primary Accession	Q9UNZ2
Other Accession	Q3SZC4, Q9CZ44, Q5RBG3, O35987
Reactivity	**Human, Mouse**
Predicted	**Bovine, Rat**
Host	**Rabbit**
Clonality	**polyclonal**
Isotype	**Rabbit IgG**
Calculated MW	**40573**

NSFL1C Antibody (Center) - Additional Information

Gene ID 55968

Other Names
NSFL1 cofactor p47, UBX domain-containing protein 2C, p97 cofactor p47, NSFL1C, UBXN2C

Target/Specificity
This NSFL1C antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 214-248 amino acids from the Central region of human NSFL1C.

Dilution
WB~~1:2000
IHC-P~~1:25
FC~~1:25

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
NSFL1C Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

NSFL1C Antibody (Center) - Protein Information

Name NSFL1C

Synonyms UBXN2C

Function Reduces the ATPase activity of VCP (By similarity). Necessary for the fragmentation of Golgi stacks during mitosis and for VCP- mediated reassembly of Golgi stacks after mitosis (By similarity). May play a role in VCP-mediated formation of transitional endoplasmic reticulum (tER) (By similarity). Inhibits the activity of CTSL (in vitro) (PubMed:15498563). Together with UBXN2B/p37, regulates the centrosomal levels of kinase AURKA/Aurora A during mitotic progression by promoting AURKA removal from centrosomes in prophase (PubMed:23649807). Also, regulates spindle orientation during mitosis (PubMed:23649807).


Cellular Location
Nucleus {ECO:0000250|UniProtKB:O35987}. Golgi apparatus, Golgi stack {ECO:0000250|UniProtKB:O35987}. Chromosome {ECO:0000250|UniProtKB:O35987}. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome {ECO:0000250|UniProtKB:O35987} Note=Predominantly nuclear in interphase cells. Bound to the axial elements of sex chromosomes in pachytene spermatocytes. A small proportion of the protein is cytoplasmic, associated with Golgi stacks Localizes to centrosome during mitotic prophase and metaphase {ECO:0000250|UniProtKB:O35987}

NSFL1C Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

NSFL1C Antibody (Center) - Images




All lanes : Anti-NSFL1C Antibody (Center) at 1:2000 dilution Lane 1: 293 whole cell lysate Lane 2: A431 whole cell lysate Lane 3: NCI-H1299 whole cell lysate Lane 4: U-251 MG whole cell lysate Lane 5: Mouse brain lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG,

(H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 41 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



AP22236c staining NSFL1C in human testis tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0. 5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hours at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.



Overlay histogram showing A431 cells stained with AP22236c(green line). The cells were fixed with 2% paraformaldehyde and then permeabilized with 90% methanol for 10 min. The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (1:25 dilution) for 60 min at 37ºC. The secondary antibody used was Goat-Anti-Rabbit IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed at 1/200 dilution for 40 min at Room temperature. Isotype control antibody (blue line) was rabbit IgG1 (1µg/1x10^6 cells) used under the same conditions. Acquisition of >10, 000 events was performed.

NSFL1C Antibody (Center) - Background

Reduces the ATPase activity of VCP. Necessary for the fragmentation of Golgi stacks during mitosis and for VCP-mediated reassembly of Golgi stacks after mitosis. May play a role in VCP-mediated formation of transitional endoplasmic reticulum (tER) (By similarity). Inhibits the activity of CTSL (in vitro).

NSFL1C Antibody (Center) - References

Yue P.,et al.Submitted (AUG-1998) to the EMBL/GenBank/DDBJ databases.
Hu R.-M.,et al.Proc. Natl. Acad. Sci. U.S.A. 97:9543-9548(2000).

Zhang Q.-H.,et al.Genome Res. 10:1546-1560(2000).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Deloukas P.,et al.Nature 414:865-871(2001).