10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# Ribophorin 2 (RPN2) Antibody (N-term)

**Purified Rabbit Polyclonal Antibody (Pab)**
**Catalog # AP2410a**

## Specification

### Ribophorin 2 (RPN2) Antibody (N-term) - Product Information

| | |
|---|---|
| Application | **IHC-P,E** |
| Primary Accession | P04844 |
| Other Accession | P25235, Q9GL01, Q3SZI6, NP_002942 |
| Reactivity | **Human** |
| Predicted | **Bovine, Pig, Rat** |
| Host | **Rabbit** |
| Clonality | **Polyclonal** |
| Isotype | **Rabbit IgG** |
| Calculated MW | **69284** |
| Antigen Region | **16-46** |

### Ribophorin 2 (RPN2) Antibody (N-term) - Additional Information

**Gene ID** 6185

**Other Names**
Dolichyl-diphosphooligosaccharide--protein glycosyltransferase subunit 2, Dolichyl-diphosphooligosaccharide--protein glycosyltransferase 63 kDa subunit, RIBIIR, Ribophorin II, RPN-II, Ribophorin-2, RPN2

**Target/Specificity**
This Ribophorin 2 (RPN2) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 16-46 amino acids from the N-terminal region of human Ribophorin 2 (RPN2).

**Dilution**
IHC-P~~1:50~100

**Format**
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

**Storage**
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**
Ribophorin 2 (RPN2) Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

### Ribophorin 2 (RPN2) Antibody (N-term) - Protein Information

**Name** RPN2 (HGNC:10382)

**Function** Subunit of the oligosaccharyl transferase (OST) complex that catalyzes the initial transfer of a defined glycan (Glc(3)Man(9)GlcNAc(2) in eukaryotes) from the lipid carrier dolichol-pyrophosphate to an asparagine residue within an Asn-X-Ser/Thr consensus motif in nascent polypeptide chains, the first step in protein N-glycosylation (PubMed:31831667). N-glycosylation occurs cotranslationally and the complex associates with the Sec61 complex at the channel-forming translocon complex that mediates protein translocation across the endoplasmic reticulum (ER). All subunits are required for a maximal enzyme activity.

**Cellular Location**
Endoplasmic reticulum {ECO:0000250|UniProtKB:F1PCT7}. Endoplasmic reticulum membrane; Multi- pass membrane protein


**Tissue Location**
Expressed in all tissues tested.

## Ribophorin 2 (RPN2) Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

## Ribophorin 2 (RPN2) Antibody (N-term) - Images




Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.

## Ribophorin 2 (RPN2) Antibody (N-term) - Background

RNP2 a type I integral membrane protein found only in the rough endoplasmic reticulum. The

encoded protein is part of an N-oligosaccharyl transferase complex that links high mannose oligosaccharides to asparagine residues found in the Asn-X-Ser/Thr consensus motif of nascent polypeptide chains. This protein is similar in sequence to the yeast oligosaccharyl transferase subunit SWP1.

**Ribophorin 2 (RPN2) Antibody (N-term) - References**

Kelleher, D.J., et al., Mol. Cell 12(1):101-111 (2003). Fu, J., et al., J. Biol. Chem. 275(6):3984-3990 (2000). Loffler, C., et al., Hum. Genet. 87(2):221-222 (1991). Crimaudo, C., et al., EMBO J. 6(1):75-82 (1987). Stoffel, M., et al., Hum. Mol. Genet. 1 (8), 656 (1992).