

CPT1A Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP2524b**Specification**

CPT1A Antibody (C-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	P50416
Reactivity	Bovine, Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	88368
Antigen Region	606-636

CPT1A Antibody (C-term) - Additional Information**Gene ID** 1374**Other Names**

Carnitine O-palmitoyltransferase 1, liver isoform, CPT1-L, Carnitine O-palmitoyltransferase I, liver isoform, CPT I, CPTI-L, Carnitine palmitoyltransferase 1A, CPT1A, CPT1

Target/Specificity

This CPT1A antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 606-636 amino acids from the C-terminal region of human CPT1A.

Dilution

WB~~1:1000
IHC-P~~1:50~100

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CPT1A Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

CPT1A Antibody (C-term) - Protein Information**Name** CPT1A ([HGNC:2328](#))**Synonyms** CPT1

Function Catalyzes the transfer of the acyl group of long-chain fatty acid-CoA conjugates onto carnitine, an essential step for the mitochondrial uptake of long-chain fatty acids and their subsequent beta-oxidation in the mitochondrion (PubMed:[9691089](#), PubMed:[11350182](#), PubMed:[14517221](#), PubMed:[16651524](#)). Plays an important role in hepatic triglyceride metabolism (By similarity).

Cellular Location

Mitochondrion outer membrane; Multi-pass membrane protein

Tissue Location


Strong expression in kidney and heart, and lower in liver and skeletal muscle

CPT1A Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CPT1A Antibody (C-term) - Images



Western blot analysis of anti-CPT1A Pab (Cat. #AP2524b) in Y79 cell line lysate (35ug/lane). CPT1A (arrow) was detected using the purified Pab.



Left image is paraformaldehyde-fixed and paraffin-embedded cow lactating with CPT1A Pab (Cat. #AP2524b), which was peroxidase-conjugated to the secondary antibody, followed by AEC staining, right image is contrast, did not add the antibody. This data was kindly offered by Hideaki Hayashi, University of Bern, Switzerland.

CPT1A Antibody (C-term) - Background

The mitochondrial oxidation of long-chain fatty acids is initiated by the sequential action of carnitine palmitoyltransferase I (which is located in the outer membrane and is detergent-labile) and carnitine palmitoyltransferase II (which is located in the inner membrane and is detergent-stable), together with a carnitine-acylcarnitine translocase. CPT I is the key enzyme in the carnitine-dependent transport across the mitochondrial inner membrane and its deficiency results in a decreased rate of fatty acid beta-oxidation.

CPT1A Antibody (C-term) - References

Rasmussen, B.B., et al., J. Clin. Invest. 110(11):1687-1693 (2002).
Ogawa, E., et al., J. Hum. Genet. 47(7):342-347 (2002).
Cook, G.A., et al., Am. J. Med. Sci. 318(1):43-48 (1999).
Ijlst, L., et al., J. Clin. Invest. 102(3):527-531 (1998).
Britton, C.H., et al., Genomics 40(1):209-211 (1997).

CPT1A Antibody (C-term) - Citations

- [Molecular adaptation in adipose tissue in response to overfeeding with a high-fat diet under sedentary conditions in South Asian and Caucasian men.](#)