

### **HNF4A Antibody (S142)**

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP2778a

## **Specification**

### **HNF4A Antibody (S142) - Product Information**

Application WB,E
Primary Accession P41235

Other Accession <u>Q91766</u>, <u>P22449</u>, <u>P49698</u>, <u>NP 849180</u>

Reactivity Human

Predicted Mouse, Rat, Xenopus

Host Rabbit
Clonality Polyclonal
Isotype Rabbit IgG
Calculated MW 52785
Antigen Region 118-146

## HNF4A Antibody (S142) - Additional Information

#### **Gene ID 3172**

### **Other Names**

Hepatocyte nuclear factor 4-alpha, HNF-4-alpha, Nuclear receptor subfamily 2 group A member 1, Transcription factor 14, TCF-14, Transcription factor HNF-4, HNF-4, HNF-4, HNF-4, TCF-14

## Target/Specificity

This HNF4A antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 118-146 amino acids from human HNF4A.

#### **Dilution**

WB~~1:1000

### **Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

#### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

### **Precautions**

HNF4A Antibody (S142) is for research use only and not for use in diagnostic or therapeutic procedures.

## HNF4A Antibody (S142) - Protein Information

### Name HNF4A



## Synonyms HNF4, NR2A1, TCF14

**Function** Transcriptional regulator which controls the expression of hepatic genes during the transition of endodermal cells to hepatic progenitor cells, facilitating the recruitment of RNA pol II to the promoters of target genes (PubMed:30597922). Activates the transcription of CYP2C38 (By similarity). Represses the CLOCK-BMAL1 transcriptional activity and is essential for circadian rhythm maintenance and period regulation in the liver and colon cells (PubMed:30530698).


**Cellular Location** Nucleus.

## **HNF4A Antibody (S142) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

# HNF4A Antibody (S142) - Images




Western blot analysis of anti-HNF4A Antibody (S142) (Cat.#AP2778a) in NCI-H460 cell line lysates (35ug/lane). HNF4A (arrow) was detected using the purified Pab.

## HNF4A Antibody (S142) - Background

HNF4A is a nuclear transcription factor which binds DNA as a homodimer. This protein controls the expression of several genes, including hepatocyte nuclear factor 1 alpha, a transcription factor which regulates the expression of several hepatic genes. This protein may play a role in development of the liver, kidney, and intestines. Mutations in this protein have been associated with monogenic autosomal dominant non-insulin-dependent diabetes mellitus type I.

### **HNF4A Antibody (S142) - References**

Sugai, M., Pathol. Int. 58 (11), 681-686 (2008)





Lee, S.S., Hepatology 48 (2), 635-645 (2008) Guo, H., Biochem. J. 394 (PT 2), 379-387 (2006) Hong, Y.H., J. Biol. Chem. 278 (30), 27495-27501 (2003)