10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

ENPP2 Antibody (Center K416)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP2854C

Specification

ENPP2 Antibody (Center K416) - Product Information

Application	**WB, IHC-P, FC,E**
Primary Accession	Q13822
Other Accession	A1A4K5
Reactivity	**Human**
Predicted	**Bovine**
Host	**Rabbit**
Clonality	**Polyclonal**
Isotype	**Rabbit IgG**
Antigen Region	**401-430**

ENPP2 Antibody (Center K416) - Additional Information

Gene ID 5168

Other Names
Ectonucleotide pyrophosphatase/phosphodiesterase family member 2, E-NPP 2, Autotaxin, Extracellular lysophospholipase D, LysoPLD, ENPP2, ATX, PDNP2

Target/Specificity
This ENPP2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 401-430 amino acids from the Central region of human ENPP2.

Dilution
WB~~1:1000
IHC-P~~1:10~50
FC~~1:10~50

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
ENPP2 Antibody (Center K416) is for research use only and not for use in diagnostic or therapeutic procedures.

ENPP2 Antibody (Center K416) - Protein Information

Name ENPP2

Function Hydrolyzes lysophospholipids to produce the signaling molecule lysophosphatidic acid (LPA) in extracellular fluids (PubMed:15769751, PubMed:26371182, PubMed:27754931, PubMed:14500380, PubMed:12354767). Major substrate is lysophosphatidylcholine (PubMed:12176993, PubMed:27754931, PubMed:14500380). Can also act on sphingosylphosphorylcholine producing sphingosine-1-phosphate, a modulator of cell motility (PubMed:14500380). Can hydrolyze, in vitro, bis-pNPP, to some extent pNP-TMP, and barely ATP (PubMed:15769751, PubMed:12176993). Involved in several motility-related processes such as angiogenesis and neurite outgrowth. Acts as an angiogenic factor by stimulating migration of smooth muscle cells and microtubule formation (PubMed:11559573). Stimulates migration of melanoma cells, probably via a pertussis toxin-sensitive G protein (PubMed:1733949). May have a role in induction of parturition (PubMed:12176993). Possible involvement in cell proliferation and adipose tissue development (Probable). Tumor cell motility-stimulating factor (PubMed:1733949, PubMed:11559573). Required for LPA production in activated platelets, cleaves the sn-1 lysophospholipids to generate sn-1 lysophosphatidic acids containing predominantly 18:2 and 20:4 fatty acids (PubMed:21393252). Shows a preference for the sn-1 to the sn-2 isomer of 1-O-alkyl-sn-glycero-3- phosphocholine (lyso-PAF) (PubMed:21393252).

Cellular Location
Secreted

Tissue Location
Detected in blood plasma (at protein level) (PubMed:12176993, PubMed:26371182). Predominantly expressed in brain, placenta, ovary, and small intestine. Expressed in a number of carcinomas such as hepatocellular and prostate carcinoma, neuroblastoma and non-small-cell lung cancer. Expressed in body fluids such as plasma, cerebral spinal fluid (CSF), saliva, follicular and amniotic fluids. Not detected in leukocytes. Isoform 1 is more highly expressed in peripheral tissues than in the central nervous system (CNS) Adipocytes only express isoform 1. Isoform 3 is more highly expressed in the brain than in peripheral tissues.


ENPP2 Antibody (Center K416) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

ENPP2 Antibody (Center K416) - Images


10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999



Western blot analysis of anti-ENPP2 Antibody (Center K416) (Cat.#AP2854c) in Y79 cell line lysates (35ug/lane).ENPP2(arrow) was detected using the purified Pab.




Western blot analysis of anti-ENPP2 Antibody (Center K416) (Cat.#AP2854c) in mouse cerebellum tissue lysates (35ug/lane).ENPP2(arrow) was detected using the purified Pab.



10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

ENPP2 Antibody (Center K416) (Cat. #AP2854C)immunohistochemistry analysis in formalin fixed and paraffin embedded human hepatocarcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining.This data demonstrates the use of ENPP2 Antibody (Center K416) for immunohistochemistry. Clinical relevance has not been evaluated.



ENPP2 Antibody (Center K416) (Cat. #AP2854c) flow cytometric analysis of MDA-MB435 cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

ENPP2 Antibody (Center K416) - Background

ENPP2 functions as both a phosphodiesterase, which cleaves phosphodiester bonds at the 5' end of oligonucleotides, and a phospholipase, which catalyzes production of lysophosphatidic acid (LPA) in extracellular fluids. LPA evokes growth factor-like responses including stimulation of cell proliferation and chemotaxis. This protein stimulates the motility of tumor cells and has angiogenic properties, and its expression is upregulated in several kinds of carcinomas. This protein is secreted and further processed to make the biologically active form.

ENPP2 Antibody (Center K416) - References

Kawagoe H., Soma O., Goji J., Nishimura N., Narita M.,Genomics 30:380-384(1995)
Nam S.W., Clair T., Kim Y.S., McMarlin A.,Cancer Res. 61:6938-6944(2001)
The MGC Project Team Genome Res. 14:2121-2127(2004)