

LSM1 Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP2863c

Specification

LSM1 Antibody (Center) - Product Information

Application	WB,E
Primary Accession Other Accession	<u>015116</u> <u>08VC85, 05E9Z8</u>
Reactivity	Human, Mouse
Predicted	Bovine
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	15179
Antigen Region	33-61

LSM1 Antibody (Center) - Additional Information

Gene ID 27257

Other Names

U6 snRNA-associated Sm-like protein LSm1, Cancer-associated Sm-like, Small nuclear ribonuclear CaSm, LSM1, CASM

Target/Specificity

This LSM1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 33-61 amino acids from the Central region of human LSM1.

Dilution WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

LSM1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

LSM1 Antibody (Center) - Protein Information

Name LSM1



Synonyms CASM

Function Plays a role in the degradation of histone mRNAs, the only eukaryotic mRNAs that are not polyadenylated (PubMed:<u>18172165</u>). Probably also part of an LSm subunits-containing complex involved in the general process of mRNA degradation (By similarity).


Cellular Location Cytoplasm. Cytoplasm, P-body

LSM1 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- <u>Dot Blot</u>
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

LSM1 Antibody (Center) - Images



Western blot analysis of anti-LSM1 Antibody (Center) (Cat.#AP2863c) in mouse kidney tissue lysates (35ug/lane). LSM1(arrow) was detected using the purified Pab.

LSM1 Antibody (Center) - Background

Sm-like proteins were identified in a variety of organisms based on sequence homology with the Sm protein family. Sm-like proteins contain the Sm sequence motif, which consists of 2 regions separated by a linker of variable length that folds as a loop. The Sm-like proteins are thought to form a stable heteromer present in tri-snRNP particles, which are important for pre-mRNA splicing.

LSM1 Antibody (Center) - References

Schweinfest C.W., Graber M.W., Chapman J.M., Papas T.S., Baron P.L., Cancer Res. 57:2961-2965(1997)