

**Phospho-ILK(T173) Antibody**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP3488a**

**Specification**

---

**Phospho-ILK(T173) Antibody - Product Information**

Application	DB,E
Primary Accession	<a href="#">Q13418</a>
Other Accession	<a href="#">Q99J82</a> , <a href="#">Q55222</a> , <a href="#">Q3SWY2</a> , <a href="#">NP_004508</a> , <a href="#">Q9DF58</a>
Reactivity	Human
Predicted	Bovine, Chicken, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG

**Phospho-ILK(T173) Antibody - Additional Information**

**Gene ID** 3611

**Other Names**

Integrin-linked protein kinase, 59 kDa serine/threonine-protein kinase, ILK-1, ILK-2, p59ILK, ILK, ILK1, ILK2

**Target/Specificity**

This ILK Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding T173 of human ILK.

**Dilution**

DB~~1:500

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

Phospho-ILK(T173) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**Phospho-ILK(T173) Antibody - Protein Information**

**Name** ILK ([HGNC:6040](#))

**Function** Receptor-proximal protein kinase regulating integrin-mediated signal transduction (PubMed:[8538749](#), PubMed:[9736715](#)). May act as a mediator of inside-out integrin signaling

(PubMed:[10712922](#)). Focal adhesion protein part of the complex ILK-PINCH (PubMed:[10712922](#)). This complex is considered to be one of the convergence points of integrin- and growth factor-signaling pathway (PubMed:[10712922](#)). Could be implicated in mediating cell architecture, adhesion to integrin substrates and anchorage-dependent growth in epithelial cells (PubMed:[10712922](#)). Regulates cell motility by forming a complex with PARVB (PubMed:[32528174](#)). Phosphorylates beta-1 and beta-3 integrin subunit on serine and threonine residues, but also AKT1 and GSK3B (PubMed:[8538749](#), PubMed:[9736715](#)).

#### Cellular Location

Cell junction, focal adhesion. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cell projection, lamellipodium {ECO:0000250|UniProtKB:O55222}. Cytoplasm, myofibril, sarcomere

#### Tissue Location


Highly expressed in heart followed by skeletal muscle, pancreas and kidney. Weakly expressed in placenta, lung and liver

### Phospho-ILK(T173) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Phospho-ILK(T173) Antibody - Images



Dot blot analysis of anti-ILK-pT173 Pab (RB13894) on nitrocellulose membrane. 50ng of Phospho-peptide or Non Phospho-peptide per dot were adsorbed. Antibody working concentrations are 0.5ug per ml.

### Phospho-ILK(T173) Antibody - Background

Transduction of extracellular matrix signals through integrins influences intracellular and extracellular functions, and appears to require interaction of integrin cytoplasmic domains with cellular proteins. Integrin-linked kinase (ILK), interacts with the cytoplasmic domain of beta-1 integrin. ILK is a serine/threonine protein kinase with 4 ankyrin-like repeats, which associates with the cytoplasmic domain of beta integrins and acts as a proximal receptor kinase regulating integrin-mediated signal transduction.

#### **Phospho-ILK(T173) Antibody - References**

Li, Y., et al., J. Clin. Invest. 112(4):503-516 (2003).  
Troussard, A.A., et al., J. Biol. Chem. 278(25):22374-22378 (2003).  
Marotta, A., et al., Br. J. Cancer 88(11):1755-1762 (2003).  
Cordes, N., et al., Br. J. Cancer 88(9):1470-1479 (2003).  
Fukuda, T., et al., J. Cell Biol. 160(7):1001-1008 (2003).

#### **Phospho-ILK(T173) Antibody - Citations**

- [Preservation of epithelial progenitor cells from collagenase-digested oral mucosa during ex vivo cultivation.](#)
- [Preservation of Human Limbal Epithelial Progenitor Cells on Carbodiimide Cross-linked Amniotic Membrane via Integrin-Linked Kinase-mediated Wnt Activation.](#)