

Phospho-SOX2(S83) Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP3651a

Specification

Phospho-SOX2(S83) Antibody - Product Information

Application DB,E
Primary Accession P48431

Other Accession P48432, P54231

Reactivity Human

Predicted Mouse, Sheep

Host Rabbit
Clonality Polyclonal
Isotype Rabbit IgG
Calculated MW 34310

Phospho-SOX2(S83) Antibody - Additional Information

Gene ID 6657

Other Names

Transcription factor SOX-2, SOX2

Target/Specificity

This SOX2 Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding S83 of human SOX2.

Dilution

DB~~1:500

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Phospho-SOX2(S83) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Phospho-SOX2(S83) Antibody - Protein Information

Name SOX2

Function Transcription factor that forms a trimeric complex with OCT4 on DNA and controls the expression of a number of genes involved in embryonic development such as YES1, FGF4, UTF1



Tel: 858.875.1900 Fax: 858.875.1999

and ZFP206 (By similarity). Binds to the proximal enhancer region of NANOG (By similarity). Critical for early embryogenesis and for embryonic stem cell pluripotency (PubMed:18035408). Downstream SRRT target that mediates the promotion of neural stem cell self-renewal (By similarity). Keeps neural cells undifferentiated by counteracting the activity of proneural proteins and suppresses neuronal differentiation (By similarity). May function as a switch in neuronal development (By similarity).

Cellular Location


Nucleus speckle {ECO:0000250|UniProtKB:Q05066}. Cytoplasm {ECO:0000250|UniProtKB:Q05738}. Nucleus {ECO:0000250|UniProtKB:Q05738}. Note=Acetylation contributes to its nuclear localization and deacetylation by HDAC3 induces a cytoplasmic delocalization (By similarity). Colocalizes in the nucleus with ZNF208 isoform KRAB-O and tyrosine hydroxylase (TH) (By similarity) Colocalizes with SOX6 in speckles. Colocalizes with CAML in the nucleus (By similarity). Nuclear import is facilitated by XPO4, a protein that usually acts as a nuclear export signal receptor (By similarity) {ECO:0000250|UniProtKB:Q05066, ECO:0000250|UniProtKB:Q05738}

Phospho-SOX2(S83) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Phospho-SOX2(S83) Antibody - Images



Dot blot analysis of anti-Phospho-SOX2-S83 Antibody (Cat. #AP3651a) on nitrocellulose membrane. 50ng of Phospho-peptide or Non Phospho-peptide per dot were adsorbed. Antibody working concentrations are 0.5ug per ml.

Phospho-SOX2(S83) Antibody - Background

SOX2 is a member of the SRY-related HMG-box (SOX) family of transcription factors involved in the





regulation of embryonic development and in the determination of cell fate. This protein may act as a transcriptional activator after forming a protein complex with other proteins. Mutations in the SOX2 gene have been associated with bilateral anophthalmia, a severe form of structural eye malformation.

Phospho-SOX2(S83) Antibody - References

Remenyi, A., et al., Genes Dev. 17(16):2048-2059 (2003). Wiebe, M.S., et al., J. Biol. Chem. 278(20):17901-17911 (2003). Fantes, J., et al., Nat. Genet. 33(4):461-463 (2003). Schepers, G.E., et al., Dev. Cell 3(2):167-170 (2002). Kamachi, Y., et al., Trends Genet. 16(4):182-187 (2000).