

**S47A2 Antibody (N-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP4722a**

**Specification**

---

**S47A2 Antibody (N-term) - Product Information**

Application	WB,E
Primary Accession	<a href="#">Q86VL8</a>
Reactivity	Rat
Predicted	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	125-153

**S47A2 Antibody (N-term) - Additional Information**

**Gene ID** 146802

**Other Names**

Multidrug and toxin extrusion protein 2, MATE-2, hMATE-2, Kidney-specific H(+)/organic cation antiporter, Solute carrier family 47 member 2, SLC47A2, MATE2

**Target/Specificity**

This S47A2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 125-153 amino acids from the N-terminal region of human S47A2.

**Dilution**

WB~~1:500

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

S47A2 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**S47A2 Antibody (N-term) - Protein Information**

**Name** SLC47A2 ([HGNC:26439](#))

**Synonyms** MATE2

**Function** Multidrug efflux pump that functions as a H(+)/organic cation antiporter. Mediates the efflux of cationic compounds, such as the model cations, tetraethylammonium (TEA) and 1-methyl-4-phenylpyridinium (MPP+), the platinum-based drug oxaliplatin or weak bases that are positively charged at physiological pH, cimetidine, the platinum-based drugs cisplatin and oxaliplatin or the antidiabetic drug metformin. Mediates the efflux of endogenous compounds such as, creatinine, thiamine and estrone-3-sulfate. Plays a physiological role in the excretion of drugs, toxins and endogenous metabolites through the kidney.

#### Cellular Location

Cell membrane; Multi-pass membrane protein. Apical cell membrane; Multi-pass membrane protein. Note=Detected in the renal urinary tubules.

#### Tissue Location


[Isoform 1]: High expression in kidney. Very small expression in adrenal gland and lung. [Isoform 6]: Ubiquitously expressed in all tissues examined except the kidney.

### S47A2 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### S47A2 Antibody (N-term) - Images



Anti-S47A2 Antibody (N-term) at 1:500 dilution + rat kidney tissue lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 65 kDa Blocking/Dilution buffer: 5% NFDm/TBST.

### S47A2 Antibody (N-term) - Background

S47A2 belongs to a family of transporters involved in excretion of toxic electrolytes, both endogenous and exogenous, through urine and bile. This transporter family shares homology with the bacterial MATE (multidrug and toxin extrusion) protein family responsible for drug resistance. This gene is one of two members of the MATE transporter family located near each other on chromosome 17.

#### **S47A2 Antibody (N-term) - References**

Toyama, K., et al. Pharmacogenet. Genomics 20(2):135-138(2010)  
Kajiwara, M., et al. J. Hum. Genet. 54(1):40-46(2009)  
Ohta, K.Y., et al. J Pharm Pharm Sci 12(3):388-396(2009)