

**FA2H Antibody (Center)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP4799c****Specification**

---

**FA2H Antibody (Center) - Product Information**

Application	WB, IHC-P,E
Primary Accession	<a href="#">Q7L5A8</a>
Other Accession	<a href="#">Q2LAM0</a> , <a href="#">Q5MPP0</a> , <a href="#">Q4R4P4</a>
Reactivity	Human, Mouse
Predicted	Monkey, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	42791
Antigen Region	121-148

**FA2H Antibody (Center) - Additional Information****Gene ID** 79152**Other Names**

Fatty acid 2-hydroxylase, 1---, Fatty acid alpha-hydroxylase, FA2H, FAAH

**Target/Specificity**

This FA2H antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 121-148 amino acids from the Central region of human FA2H.

**Dilution**

WB~~1:1000

IHC-P~~1:50~100

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

FA2H Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

**FA2H Antibody (Center) - Protein Information****Name** FA2H

**Function** Catalyzes the hydroxylation of free fatty acids at the C-2 position to produce 2-hydroxy fatty acids, which are building blocks of sphingolipids and glycosphingolipids common in neural tissue and epidermis (PubMed:[15337768](#), PubMed:[15863841](#), PubMed:[17355976](#), PubMed:[22517924](#)). FA2H is stereospecific for the production of (R)-2- hydroxy fatty acids (PubMed:[22517924](#)). Plays an essential role in the synthesis of galactosphingolipids of the myelin sheath (By similarity). Responsible for the synthesis of sphingolipids and glycosphingolipids involved in the formation of epidermal lamellar bodies critical for skin permeability barrier (PubMed:[17355976](#)). Participates in the synthesis of glycosphingolipids and a fraction of type II wax diesters in sebaceous gland, specifically regulating hair follicle homeostasis (By similarity). Involved in the synthesis of sphingolipids of plasma membrane rafts, controlling lipid raft mobility and trafficking of raft-associated proteins (By similarity).

#### Cellular Location

Endoplasmic reticulum membrane {ECO:0000250|UniProtKB:Q5MPP0}; Multi-pass membrane protein. Microsome membrane; Multi-pass membrane protein

#### Tissue Location


Detected in differentiating cultured keratinocytes (at protein level). Detected in epidermis and cultured keratinocytes (PubMed:[17355976](#)). Highly expressed in brain and colon. Detected at lower levels in testis, prostate, pancreas and kidney (PubMed:[15337768](#)).

#### FA2H Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


#### FA2H Antibody (Center) - Images



Western blot analysis of FA2H Antibody (Center) (Cat. #AP4799c) in ZR-75-1 cell line lysates (35ug/lane). FA2H (arrow) was detected using the purified Pab.



Western blot analysis of FA2H Antibody (Center) (Cat. #AP4799c) in mouse brain tissue lysates (35ug/lane). FA2H (arrow) was detected using the purified Pab.



FA2H Antibody (Center) (Cat. #AP4799c) IHC analysis in formalin fixed and paraffin embedded brain tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the FA2H Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.

### **FA2H Antibody (Center) - Background**

FA2H is a protein that catalyzes the synthesis of 2-hydroxysphingolipids, a subset of sphingolipids that contain 2-hydroxy fatty acids. Sphingolipids play roles in many cellular processes and their structural diversity arises from modification of the hydrophobic ceramide moiety, such as by 2-hydroxylation of the N-acyl chain, and the existence of many different head groups.

### **FA2H Antibody (Center) - References**

- Wheeler, H.E., et al. PLoS Genet. 5 (10), E1000685 (2009)
- Edvardson, S., et al. Am. J. Hum. Genet. 83(5):643-648(2008)
- Uchida, Y., et al. J. Biol. Chem. 282(18):13211-13219(2007)