

**C19orf21 Antibody (N-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP4851a****Specification**

---

**C19orf21 Antibody (N-term) - Product Information**

Application	WB, IHC-P, FC,E
Primary Accession	<a href="#">Q8IVT2</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	75357
Antigen Region	63-91

**C19orf21 Antibody (N-term) - Additional Information****Gene ID** 126353**Other Names**

Mitotic interactor and substrate of PLK1, Mitotic spindle positioning protein, MISP, C19orf21

**Target/Specificity**

This C19orf21 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 63-91 amino acids from the N-terminal region of human C19orf21.

**Dilution**

WB~~1:1000  
IHC-P~~1:50~100  
FC~~1:10~50

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

C19orf21 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**C19orf21 Antibody (N-term) - Protein Information****Name** MISP ([HGNC:27000](#))**Function** Plays a role in mitotic spindle orientation and mitotic progression. Regulates the

distribution of dynactin at the cell cortex in a PLK1-dependent manner, thus stabilizing cortical and astral microtubule attachments required for proper mitotic spindle positioning. May link microtubules to the actin cytoskeleton and focal adhesions. May be required for directed cell migration and centrosome orientation. May also be necessary for proper stacking of the Golgi apparatus.

#### Cellular Location

Cell junction, focal adhesion. Cytoplasm, cytoskeleton. Cytoplasm, cell cortex.


Note=Predominantly localizes to cortical actin structures during interphase and mitosis. Present in retraction fibers, which are formed at former adhesion sites during mitosis, and at spicular membrane protrusions in re-attaching cytokinetic cells. Partially colocalizes with cytoplasmic F-actin. Not detected at microtubules at interphase, nor at spindle during mitosis

#### C19orf21 Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


#### C19orf21 Antibody (N-term) - Images



Western blot analysis of C19orf21 Antibody (N-term) (Cat. #AP4851a) in HepG2 cell line lysates (35ug/lane). C19orf21 (arrow) was detected using the purified Pab.



C19orf21 Antibody (N-term) (Cat. #AP4851a) IHC analysis in formalin fixed and paraffin embedded human cervix carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the C19orf21 Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.



C19orf21 Antibody (N-term) (Cat. #AP4851a) flow cytometric analysis of HepG2 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

### **C19orf21 Antibody (N-term) - Background**

The exact function of C19orf21 remains unknown.

### **C19orf21 Antibody (N-term) - References**

- Olsen, J.V., et al. Cell 127(3):635-648(2006)
- Nousiainen, M., et al. Proc. Natl. Acad. Sci. U.S.A. 103(14):5391-5396(2006)
- Kim, J.E., et al. J. Proteome Res. 4(4):1339-1346(2005)