10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Cyclin D2 Antibody (Ab-280)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP50014

Specification

Cyclin D2 Antibody (Ab-280) - Product Information

Application	**WB**
Primary Accession	P30279
Reactivity	**Human, Mouse, Rat**
Host	**Rabbit**
Clonality	**Polyclonal**
Calculated MW	**33 KDa**
Antigen Region	**253-285**

Cyclin D2 Antibody (Ab-280) - Additional Information

Gene ID 894

Other Names
G1/S-specific cyclin-D2, CCND2

Dilution
WB~~ 1:500-1:1000

Format
Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.09% (W/V) sodium azide and 50% glycerol.

Storage Conditions
-20℃

Cyclin D2 Antibody (Ab-280) - Protein Information

Name CCND2 {ECO:0000303|PubMed:1386336, ECO:0000312|HGNC:HGNC:1583}

Function
Regulatory component of the cyclin D2-CDK4 (DC) complex that phosphorylates and inhibits members of the retinoblastoma (RB) protein family including RB1 and regulates the cell-cycle during G(1)/S transition (PubMed:8114739, PubMed:18827403). Phosphorylation of RB1 allows dissociation of the transcription factor E2F from the RB/E2F complex and the subsequent transcription of E2F target genes which are responsible for the progression through the G(1) phase (PubMed:8114739, PubMed:18827403). Hypophosphorylates RB1 in early G(1) phase (PubMed:8114739, PubMed:18827403). Cyclin

D-CDK4 complexes are major integrators of various mitogenenic and antimitogenic signals (PubMed:8114739, PubMed:18827403).


Cellular Location
Nucleus. Cytoplasm. Nucleus membrane. Note=Cyclin D-CDK4 complexes accumulate at the nuclear membrane and are then translocated into the nucleus through interaction with KIP/CIP family members

Cyclin D2 Antibody (Ab-280) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Cyclin D2 Antibody (Ab-280) - Images



Western blot analysis of lysate from RD cell line,using Cyclin D2 Antibody (Ab-280)(B8336). B8336 was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody.Lysate at 35ug.

Cyclin D2 Antibody (Ab-280) - Background

Regulatory component of the cyclin D2-CDK4 (DC) complex that phosphorylates and inhibits members of the retinoblastoma (RB) protein family including RB1 and regulates the cell-cycle during G(1)/S transition. Phosphorylation of RB1 allows dissociation of the transcription factor E2F from the RB/E2F complex and the subsequent transcription of E2F target genes which are responsible for the progression through the G(1) phase. Hypophosphorylates RB1 in early G(1) phase. Cyclin D-CDK4 complexes are major integrators of various mitogenenic and antimitogenic signals. Also substrate for SMAD3, phosphorylating SMAD3 in a cell-cycle-dependent manner and

repressing its transcriptional activity. Component of the ternary complex, cyclin D2/CDK4/CDKN1B, required for nuclear translocation and activity of the cyclin D-CDK4 complex (By similarity).

Cyclin D2 Antibody (Ab-280) - References

Xiong Y.,et al.Genomics 13:575-584(1992).
Palmero I.,et al.Oncogene 8:1049-1054(1993).
Miyajima N.,et al.Submitted (MAR-1993) to the EMBL/GenBank/DDBJ databases.
Kalnine N.,et al.Submitted (OCT-2004) to the EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).