

TAIP-2 Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP50827**Specification**

TAIP-2 Antibody - Product Information

Application	WB
Primary Accession	Q8WYN3
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	65,58 KDa
Antigen Region	180-211

TAIP-2 Antibody - Additional Information**Gene ID** 80034**Other Names**

Cysteine/serine-rich nuclear protein 3, CSRNP-3, Protein FAM130A2, TGF-beta-induced apoptosis protein 2, TAIP-2, CSRNP3, FAM130A2, TAIP2

Dilution

WB~~ 1:1000

FormatRabbit IgG in phosphate buffered saline (without Mg²⁺ and Ca²⁺), pH 7.4, 150mM NaCl, 0.09% (W/V) sodium azide and 50% glycerol.**Storage Conditions**

-20°C

TAIP-2 Antibody - Protein Information**Name** CSRNP3**Synonyms** FAM130A2, TAIP2**Function**

Binds to the consensus sequence 5'-AGAGTG-3' and has transcriptional activator activity. Plays a role in apoptosis (By similarity).

Cellular Location


Nucleus.

TAIP-2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TAIP-2 Antibody - Images



Western blot analysis of lysate from PC-3 cell line, using TAIP-2 Antibody, was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35ug.

TAIP-2 Antibody - Background

Binds to the consensus sequence 5'-AGAGTG-3' and has transcriptional activator activity. Plays a role in apoptosis (By similarity).

TAIP-2 Antibody - References

Akiyama N., et al. Submitted (JUN-2001) to the EMBL/GenBank/DDBJ databases.
Ota T., et al. Nat. Genet. 36:40-45(2004).
Hillier L.W., et al. Nature 434:724-731(2005).
Mural R.J., et al. Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.
Sjoebom T., et al. Science 314:268-274(2006).