

HDAC9 Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP50939**Specification**

HDAC9 Antibody - Product Information

Application	WB
Primary Accession	Q9UKV0
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	117,112,118 KDa
Antigen Region	1032-1061

HDAC9 Antibody - Additional Information**Gene ID** 9734**Other Names**

Histone deacetylase 9, HD9, Histone deacetylase 7B, HD7, HD7b, Histone deacetylase-related protein, MEF2-interacting transcription repressor MITR, HDAC9, HDAC7, HDAC7B, HDRP, KIAA0744, MITR

Dilution

WB~~ 1:500

Format

Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.09% (W/V) sodium azide and 54% glycerol

Storage

Store at -20 °C.Stable for 12 months from date of receipt

HDAC9 Antibody - Protein Information**Name** HDAC9**Synonyms** HDAC7, HDAC7B, HDRP, KIAA0744, MITR**Function**

Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Represses MEF2-dependent transcription.

Cellular Location

Nucleus.


Tissue Location

Broadly expressed, with highest levels in brain, heart, muscle and testis. Isoform 3 is present in human bladder carcinoma cells (at protein level).

HDAC9 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

HDAC9 Antibody - Images

Western blot analysis of lysates from HeLa, K562 cell line (from left to right), using HDAC9 Antibody (A1-0639). A1-0639 was diluted at 1:500 at each lane. A goat anti-rabbit IgG H&L (HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 20 µg per lane.