

Catenin alpha 1/2 Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP51125

Specification

Catenin alpha 1/2 Antibody - Product Information

Application Primary Accession Reactivity Host Clonality Calculated MW Antigen Region WB <u>P35221</u> Human, Mouse, Rat Rabbit Polyclonal 102 KDa 841 - 900

Catenin alpha 1/2 Antibody - Additional Information

Gene ID 1495

Other Names Catenin alpha-1, Alpha E-catenin, Cadherin-associated protein, Renal carcinoma antigen NY-REN-13, CTNNA1

Target/Specificity KLH conjugated synthetic peptide derived from human Catenin alpha 1/2

Dilution WB~~ 1:1000

Format 0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

Storage Store at -20 °C.Stable for 12 months from date of receipt

Catenin alpha 1/2 Antibody - Protein Information

Name CTNNA1

Function

Associates with the cytoplasmic domain of a variety of cadherins. The association of catenins to cadherins produces a complex which is linked to the actin filament network, and which seems to be of primary importance for cadherins cell-adhesion properties. Can associate with both E- and N-cadherins. Originally believed to be a stable component of E-cadherin/catenin adhesion complexes and to mediate the linkage of cadherins to the actin cytoskeleton at adherens junctions. In contrast, cortical actin was found to be much more dynamic than E-cadherin/catenin complexes and CTNNA1 was shown not to bind to F-actin when assembled in the complex suggesting a different linkage between actin and adherens junctions components. The homodimeric form may regulate actin filament assembly and inhibit actin branching by competing



with the Arp2/3 complex for binding to actin filaments. Involved in the regulation of WWTR1/TAZ, YAP1 and TGFB1- dependent SMAD2 and SMAD3 nuclear accumulation (By similarity). May play a crucial role in cell differentiation.

Cellular Location

[Isoform 1]: Cytoplasm, cytoskeleton. Cell junction, adherens junction. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cell junction. Note=Found at cell-cell boundaries and probably at cell-matrix boundaries

Tissue Location


Expressed ubiquitously in normal tissues.

Catenin alpha 1/2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Catenin alpha 1/2 Antibody - Images



All lanes : Anti-Catenin alpha 1/2 Antibody at 1:1000 dilution Lane 1: Hela whole cell lysates Lane 2: SH-SY5Y whole cell lysates Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L),Peroxidase conjugated at 1/10000 dilution Predicted band size : 100 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Catenin alpha 1/2 Antibody - Background

Associates with the cytoplasmic domain of a variety of cadherins. The association of catenins to cadherins produces a complex which is linked to the actin filament network, and which seems to be of primary importance for cadherins cell-adhesion properties. Can associate with both E- and



N-cadherins. Originally believed to be a stable component of E-cadherin/catenin adhesion complexes and to mediate the linkage of cadherins to the actin cytoskeleton at adherens junctions. In contrast, cortical actin was found to be much more dynamic than E-cadherin/catenin complexes and CTNNA1 was shown not to bind to F-actin when assembled in the complex suggesting a different linkage between actin and adherens junctions components. The homodimeric form may regulate actin filament assembly and inhibit actin branching by competing with the Arp2/3 complex for binding to actin filaments. May play a crucial role in cell differentiation.

Catenin alpha 1/2 Antibody - References

Furukawa Y.,et al.Cytogenet. Cell Genet. 65:74-78(1994). Oda T.,et al.Biochem. Biophys. Res. Commun. 193:897-904(1993). Rimm D.L.,et al.Biochem. Biophys. Res. Commun. 203:1691-1699(1994). Kask M.,et al.Biochem. Biophys. Res. Commun. 411:56-61(2011). Nollet F.H.,et al.Submitted (OCT-1998) to the EMBL/GenBank/DDBJ databases.