

### **GAB2 Antibody**

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP51222

## **Specification**

# **GAB2 Antibody - Product Information**

Application WB
Primary Accession O9UOC2
Reactivity Human
Host Rabbit
Clonality Polyclonal
Calculated MW 74 KDa
Antigen Region 221 - 280

# **GAB2 Antibody - Additional Information**

#### **Gene ID 9846**

#### **Other Names**

GRB2-associated-binding protein 2, GRB2-associated binder 2, Growth factor receptor bound protein 2-associated protein 2, pp100, GAB2, KIAA0571

# Target/Specificity

KLH conjugated synthetic peptide derived from human GAB2

### **Dilution**

WB~~ 1:1000

#### **Format**

0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

# **Storage**

Store at -20 °C. Stable for 12 months from date of receipt

# **GAB2 Antibody - Protein Information**

### Name GAB2

## Synonyms KIAA0571

### **Function**

Adapter protein which acts downstream of several membrane receptors including cytokine, antigen, hormone, cell matrix and growth factor receptors to regulate multiple signaling pathways. Regulates osteoclast differentiation mediating the TNFRSF11A/RANK signaling. In allergic response, it plays a role in mast cells activation and degranulation through PI-3-kinase regulation. Also involved in the regulation of cell proliferation and hematopoiesis.

# **Cellular Location**




Cytoplasm. Cell membrane. Membrane raft {ECO:0000250|UniProtKB:Q9Z1S8}

# **GAB2 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

## GAB2 Antibody - Images



Anti-GAB2 Antibodyat 1:1000 dilution + MCF-7 whole cell lysates Lysates/proteins at 20  $\mu$ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L),Peroxidase conjugated at 1/10000 dilution Predicted band size : 74 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

### **GAB2 Antibody - Background**

Adapter protein which acts downstream of several membrane receptors including cytokine, antigen, hormone, cell matrix and growth factor receptors to regulate multiple signaling pathways. Regulates osteoclast differentiation mediating the TNFRSF11A/RANK signaling. In allergic response, it plays a role in mast cells activation and degranulation through PI-3-kinase regulation. Also involved in the regulation of cell proliferation and hematopoiesis.

# **GAB2 Antibody - References**

Nishida K., et al. Blood 93:1809-1816(1999).

Nagase T., et al. DNA Res. 5:31-39(1998).

Nagase T., et al. Submitted (JAN-2004) to the EMBL/GenBank/DDBJ databases.

Taylor T.D., et al. Nature 440:497-500(2006).

Mural R.J., et al. Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.