

HSD17B2 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP5125C**Specification**

HSD17B2 Antibody (Center) - Product Information

Application	WB, IHC-P, FC,E
Primary Accession	P37059
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	42785
Antigen Region	265-294

HSD17B2 Antibody (Center) - Additional Information**Gene ID** 3294**Other Names**

Estradiol 17-beta-dehydrogenase 2, 17-beta-hydroxysteroid dehydrogenase type 2, 17-beta-HSD 2, 20 alpha-hydroxysteroid dehydrogenase, 20-alpha-HSD, E2DH, Microsomal 17-beta-hydroxysteroid dehydrogenase, Testosterone 17-beta-dehydrogenase, HSD17B2, EDH17B2

Target/Specificity

This HSD17B2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 265-294 amino acids from the Central region of human HSD17B2.

Dilution

WB~~1:1000
IHC-P~~1:10~50
FC~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

HSD17B2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

HSD17B2 Antibody (Center) - Protein Information

Name HSD17B2 ([HGNC:5211](#))

Synonyms EDH17B2, SDR9C2

Function Catalyzes the NAD-dependent oxidation of the highly active 17beta-hydroxysteroids, such as estradiol (E2), testosterone (T), and dihydrotestosterone (DHT), to their less active forms and thus regulates the biological potency of these steroids. Oxidizes estradiol to estrone, testosterone to androstenedione, and dihydrotestosterone to 5alpha-androstan-3,17-dione. Also has 20-alpha-HSD activity.

Cellular Location

Endoplasmic reticulum membrane; Single-pass type II membrane protein

Tissue Location


Expressed in placenta.

HSD17B2 Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


HSD17B2 Antibody (Center) - Images



Anti-HSD17B2 Antibody (Center) at 1:1000 dilution + MDA-MB-453 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 43 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



HSD17B2 Antibody (Center) (Cat. #AP5125c) IHC analysis in formalin fixed and paraffin embedded human prostate carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the HSD17B2 Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.



HSD17B2 Antibody (Center) (Cat. #AP5125c) flow cytometric analysis of MDA-MB435 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

HSD17B2 Antibody (Center) - Background

HSD17B2 is capable of catalyzing the interconversion of testosterone and androstenedione, as well as estradiol and estrone. HSD17B2 also has 20-alpha-HSD activity. HSD17B2 uses NADH while EDH17B3 uses NADPH.

HSD17B2 Antibody (Center) - References

- Shen, Z., et al. Endocrinology 150(11):4941-4949(2009)
- Bhavani, V., et al. Cancer Biomark 5(4):207-213(2009)
- Olsen, J.V., et al. Cell 127(3):635-648(2006)