

5-HT4 Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP51275

Specification

5-HT4 Antibody - Product Information

Application WB
Primary Accession O13639
Reactivity Human
Host Rabbit
Clonality Polyclonal
Calculated MW 43 KDa
Antigen Region 11 - 70

5-HT4 Antibody - Additional Information

Gene ID 3360

Other Names

5-hydroxytryptamine receptor 4, 5-HT-4, 5-HT4, Serotonin receptor 4, HTR4

Target/Specificity

KLH conjugated synthetic peptide derived from human 5-HT4

Dilution

WB~~ 1:1000

Format

0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

Storage

Store at -20 °C. Stable for 12 months from date of receipt

5-HT4 Antibody - Protein Information

Name HTR4

Function

This is one of the several different receptors for 5- hydroxytryptamine (serotonin), a biogenic hormone that functions as a neurotransmitter, a hormone, and a mitogen. The activity of this receptor is mediated by G proteins that stimulate adenylate cyclase.

Cellular Location

Cell membrane; Multi-pass membrane protein. Endosome. Note=Interaction with SNX27 mediates recruitment to early endosomes, while interaction with NHERF1 and EZR might target the protein to specialized subcellular regions, such as microvilli

Tissue Location




Isoform 5-HT4(A) is expressed in ileum, brain, and atrium, but not in the ventricle.

5-HT4 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

5-HT4 Antibody - Images



Anti-5-HT4 Antibodyat 1:1000 dilution + Jurkat whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L),Peroxidase conjugated at 1/10000 dilution Predicted band size: 44,41,43,45,48,47 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

5-HT4 Antibody - Background

This is one of the several different receptors for 5- hydroxytryptamine (serotonin), a biogenic hormone that functions as a neurotransmitter, a hormone, and a mitogen. The activity of this receptor is mediated by G proteins that stimulate adenylate cyclase.

5-HT4 Antibody - References

Blondel O., et al.J. Neurochem. 70:2252-2261(1998).

Van den Wyngaert I., et al. Submitted (JUN-1997) to the EMBL/GenBank/DDBJ databases.

Claeysen S., et al. NeuroReport 8:3189-3195(1997).

Syversveen T., et al. Submitted (DEC-1998) to the EMBL/GenBank/DDBJ databases.

Claeysen S., et al. Mol. Pharmacol. 55:910-920(1999).