10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

p107 Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP51467

Specification

p107 Antibody - Product Information

Application	**WB**
Primary Accession	P28749
Reactivity	**Human, Mouse, Rat**
Host	**Rabbit**
Clonality	**Polyclonal**
Calculated MW	**121 KDa**
Antigen Region	**331 - 390**

p107 Antibody - Additional Information

Gene ID 5933

Other Names
Retinoblastoma-like protein 1, 107 kDa retinoblastoma-associated protein, p107, pRb1, RBL1

Target/Specificity
KLH conjugated synthetic peptide derived from human p107

Dilution
WB~~ 1:1000

Format
0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

Storage
Store at -20 °C.Stable for 12 months from date of receipt

p107 Antibody - Protein Information

Name RBL1

Function
Key regulator of entry into cell division (PubMed:17671431). Directly involved in heterochromatin formation by maintaining overall chromatin structure and, in particular, that of constitutive heterochromatin by stabilizing histone methylation (By similarity). Recruits and targets histone methyltransferases KMT5B and KMT5C, leading to epigenetic transcriptional repression (By similarity). Controls histone H4 'Lys-20' trimethylation (By similarity). Probably acts as a transcription repressor by recruiting chromatin-modifying enzymes to promoters (By similarity). Potent inhibitor of E2F-mediated trans-activation (PubMed:8319904). May act as a tumor suppressor (PubMed:<a href="http://www.uniprot.org/citations/8319904"

target="_blank">8319904).


Cellular Location
Nucleus.

p107 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

p107 Antibody - Images



Anti-p107 Antibodyat 1:1000 dilution + 293T whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L),Peroxidase conjugated at 1/10000 dilution Predicted band size : 121 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

p107 Antibody - Background

Key regulator of entry into cell division. Directly involved in heterochromatin formation by maintaining overall chromatin structure and, in particular, that of constitutive heterochromatin by stabilizing histone methylation. Recruits and targets histone methyltransferases SUV420H1 and SUV420H2, leading to epigenetic transcriptional repression. Controls histone H4 'Lys-20' trimethylation. Probably acts as a transcription repressor by recruiting chromatin-modifying enzymes to promoters. Potent inhibitor of E2F-mediated trans-activation. Forms a complex with adenovirus E1A and with SV40 large T antigen. May bind and modulate functionally certain cellular proteins with which T and E1A compete for pocket binding. May act as a tumor suppressor.

p107 Antibody - References

Zhu L.,et al.Genes Dev. 7:1111-1125(1993).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Deloukas P.,et al.Nature 414:865-871(2001).
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.
Ewen M.E.,et al.Cell 66:1155-1164(1991).