10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

ZIP7 Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP51886

Specification

ZIP7 Antibody - Product Information

Application	**WB**
Primary Accession	Q92504
Reactivity	**Human, Mouse, Rat**
Host	**Rabbit**
Clonality	**Polyclonal**
Calculated MW	**50 KDa**
Antigen Region	**121 - 180**

ZIP7 Antibody - Additional Information

Gene ID 7922

Other Names
Zinc transporter SLC39A7, Histidine-rich membrane protein Ke4, Really interesting new gene 5 protein, Solute carrier family 39 member 7, Zrt-, Irt-like protein 7, ZIP7, SLC39A7, HKE4, RING5

Target/Specificity
KLH conjugated synthetic peptide derived from human ZIP7

Dilution
WB~~ 1:4000

Format
0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

Storage
Store at -20 °C.Stable for 12 months from date of receipt

ZIP7 Antibody - Protein Information

Name SLC39A7

Synonyms HKE4 {ECO:0000303|PubMed:14525538}, RING

Function
Transports Zn(2+) from the endoplasmic reticulum (ER)/Golgi apparatus to the cytosol, playing an essential role in the regulation of cytosolic zinc levels (PubMed:14525538, PubMed:15705588, PubMed:29980658, PubMed:28205653). Acts as a gatekeeper of zinc release from intracellular stores, requiring post-translational activation by

phosphorylation, resulting in activation of multiple downstream pathways leading to cell growth and proliferation (PubMed:29980658, PubMed:22317921, PubMed:28205653). Has an essential role in B cell development and is required for proper B cell receptor signaling (PubMed:30718914). Plays an important role in maintaining intestinal epithelial homeostasis and skin dermis development by regulating ER function (By similarity). Controls cell signaling pathways involved in glucose metabolism in skeletal muscle (By similarity). Has a protective role against ER stress in different biological contexts (PubMed:29980658, PubMed:30237509). Mediates Zn(2+)-induced ferroptosis (PubMed:33608508).

Cellular Location
Endoplasmic reticulum membrane; Multi-pass membrane protein. Golgi apparatus, cis-Golgi network membrane; Multi-pass membrane protein


Tissue Location
Widely expressed.

ZIP7 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

ZIP7 Antibody - Images



10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-ZIP7 Antibodyat 1:4000 dilution + HepG2 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L),Peroxidase conjugated at 1/10000 dilution Predicted band size : 50 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

ZIP7 Antibody - Background

Zinc transporter, that transports Zn(2+) from the endoplasmic reticulum/Golgi apparatus to the cytosol. Transport is stimulated by growth factors, such as EGF, and Ca(2+), as well as by exogenous Zn(2+).

ZIP7 Antibody - References

Ando A.,et al.Genomics 35:600-602(1996).
Vergara A.,et al.Submitted (DEC-1998) to the EMBL/GenBank/DDBJ databases.
Mungall A.J.,et al.Nature 425:805-811(2003).
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.
Taylor K.M.,et al.Biochem. J. 377:131-139(2004).