10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Rabbit Anti-MTSS1 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP52041

Specification

Rabbit Anti-MTSS1 Polyclonal Antibody - Product Information

Application	**WB**
Primary Accession	O43312
Reactivity	**Human, Mouse, Rat**
Host	**Rabbit**
Clonality	**Polyclonal**
Antigen Region	**101-200/755aa**

Rabbit Anti-MTSS1 Polyclonal Antibody - Additional Information

Gene ID 9788

Other Names
MIM; MIMA; MIMB; Metastasis suppressor protein 1; Metastasis suppressor YGL-1; Missing in metastasis protein; MTSS1; KIAA429

Dilution
WB~~1:100~1:500

Format
0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage
Store at -20 ℃ for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ℃.

Rabbit Anti-MTSS1 Polyclonal Antibody - Protein Information

Name MTSS1 (HGNC:20443)

Synonyms KIAA0429, MIM

Function
May be related to cancer progression or tumor metastasis in a variety of organ sites, most likely through an interaction with the actin cytoskeleton.

Cellular Location
Cytoplasm, cytoskeleton.

Tissue Location
Expressed in many tissues, including spleen, thymus, prostate, testis, uterus, colon, and peripheral blood


10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Rabbit Anti-MTSS1 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Rabbit Anti-MTSS1 Polyclonal Antibody - Images



Mouse testicular lysate probed with Rabbit Anti-MTSS1 Polyclonal Antibody (AP52041) at 1:300 overnight in 4˚C. Followed by conjugation to the secondary antibody at 1:5000 90min in 37˚C

Rabbit Anti-MTSS1 Polyclonal Antibody - Background

May be related to cancer progression or tumor metastasis in a variety of organ sites, most likely through an interaction with the actin cytoskeleton.