

BLNK (Phospho-Tyr84) Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP52538**Specification**

BLNK (Phospho-Tyr84) Antibody - Product Information

Application	WB, IHC
Primary Accession	Q8WV28
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	50466

BLNK (Phospho-Tyr84) Antibody - Additional Information**Gene ID** 29760**Other Names**

B-cell linker protein, B-cell adapter containing a SH2 domain protein, B-cell adapter containing a Src homology 2 domain protein, Cytoplasmic adapter protein, Src homology 2 domain-containing leukocyte protein of 65 kDa, SLP-65, BLNK, BASH, SLP65

Dilution

WB~~1:1000

IHC~~1:50~100

Format

Rabbit IgG in phosphate buffered saline (without Mg²⁺ and Ca²⁺), pH 7.4, 150mM NaCl, 0.09% (W/V) sodium azide and 50% glycerol.

Storage Conditions

-20°C

BLNK (Phospho-Tyr84) Antibody - Protein Information**Name** BLNK**Synonyms** BASH, SLP65**Function**

Functions as a central linker protein, downstream of the B- cell receptor (BCR), bridging the SYK kinase to a multitude of signaling pathways and regulating biological outcomes of B-cell function and development. Plays a role in the activation of ERK/EPHB2, MAP kinase p38 and JNK. Modulates AP1 activation. Important for the activation of NF-kappa-B and NFAT. Plays an important role in BCR- mediated PLCG1 and PLCG2 activation and Ca(2+) mobilization and is required for trafficking of the BCR to late endosomes. However, does not seem to be required for pre-BCR-mediated activation of MAP kinase and phosphatidyl-inositol 3 (PI3) kinase signaling. May be required for the RAC1-JNK pathway. Plays a critical role in orchestrating the pro-B cell to pre-B cell transition. May

play an important role in BCR- induced B-cell apoptosis.

Cellular Location

Cytoplasm. Cell membrane. Note=BCR activation results in the translocation to membrane fraction


Tissue Location

Expressed in B-cell lineage and fibroblast cell lines (at protein level). Highest levels of expression in the spleen, with lower levels in the liver, kidney, pancreas, small intestines and colon


BLNK (Phospho-Tyr84) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

BLNK (Phospho-Tyr84) Antibody - Images

Western blot analysis of extracts from K562 cells, treated with starved (24hours), using BLNK (Phospho-Tyr84) antibody.



Immunohistochemistry analysis of paraffin-embedded human brain tissue using BLNK

(Phospho-Tyr84) antibody.

BLNK (Phospho-Tyr84) Antibody - Background

Functions as a central linker protein, downstream of the B-cell receptor (BCR), bridging the SYK kinase to a multitude of signaling pathways and regulating biological outcomes of B-cell function and development. Plays a role in the activation of ERK/EPHB2, MAP kinase p38 and JNK. Modulates AP1 activation. Important for the activation of NF-kappa-B and NFAT. Plays an important role in BCR-mediated PLCG1 and PLCG2 activation and Ca(2+) mobilization and is required for trafficking of the BCR to late endosomes. However, does not seem to be required for pre-BCR- mediated activation of MAP kinase and phosphatidyl-inositol 3 (PI3) kinase signaling. May be required for the RAC1-JNK pathway. Plays a critical role in orchestrating the pro-B cell to pre-B cell transition. May play an important role in BCR-induced B-cell apoptosis.

BLNK (Phospho-Tyr84) Antibody - References

Fu C.,et al.Immunity 9:93-103(1998).
Minegishi Y.,et al.Science 286:1954-1957(1999).
Sprangers M.,et al.Oncogene 25:5180-5186(2006).
Deloukas P.,et al.Nature 429:375-381(2004).
Fu C.,et al.J. Biol. Chem. 272:27362-27368(1997).