

p53 (C-terminus) Antibody
Purified Mouse Monoclonal Antibody (Mab)
Catalog # AP52665

Specification

p53 (C-terminus) Antibody - Product Information

Application	WB
Primary Accession	P04637
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG2b
Calculated MW	53 KDa

p53 (C-terminus) Antibody - Additional Information

Gene ID 7157

Other Names

Antigen NY-CO-13;BCC7;Cellular tumor antigen p53;Cys 51 Stop;FLJ92943;HGNC11998;LFS1; p53;p53 Cellular Tumor Antigen;p53 tumor suppressor;P53_HUMAN;Phosphoprotein p53;Tp53; Transformation related protein 53;TRP53;Tumor protein p53;Tumor suppressor p53;Tumour Protein p53.

Dilution

WB~~1:1000

Format

Purified mouse monoclonal in PBS(pH 7.4) containing with 0.09% (W/V) sodium azide and 50% glycerol.

Storage

Store at -20 °C.Stable for 12 months from date of receipt

p53 (C-terminus) Antibody - Protein Information

Name TP53

Synonyms P53

Function

Acts as a tumor suppressor in many tumor types; induces growth arrest or apoptosis depending on the physiological circumstances and cell type (PubMed:11025664, PubMed:12524540, PubMed:12810724, PubMed:15186775, PubMed:15340061, PubMed:<a

[17317671](http://www.uniprot.org/citations/17317671) PubMed: [17349958](http://www.uniprot.org/citations/17349958) PubMed: [19556538](http://www.uniprot.org/citations/19556538) PubMed: [20673990](http://www.uniprot.org/citations/20673990) PubMed: [20959462](http://www.uniprot.org/citations/20959462) PubMed: [22726440](http://www.uniprot.org/citations/22726440) PubMed: [24051492](http://www.uniprot.org/citations/24051492) PubMed: [9840937](http://www.uniprot.org/citations/9840937) PubMed: [24652652](http://www.uniprot.org/citations/24652652)). Involved in cell cycle regulation as a trans-activator that acts to negatively regulate cell division by controlling a set of genes required for this process (PubMed: [11025664](http://www.uniprot.org/citations/11025664) PubMed: [12524540](http://www.uniprot.org/citations/12524540) PubMed: [12810724](http://www.uniprot.org/citations/12810724) PubMed: [15186775](http://www.uniprot.org/citations/15186775) PubMed: [15340061](http://www.uniprot.org/citations/15340061) PubMed: [17317671](http://www.uniprot.org/citations/17317671) PubMed: [17349958](http://www.uniprot.org/citations/17349958) PubMed: [19556538](http://www.uniprot.org/citations/19556538) PubMed: [20673990](http://www.uniprot.org/citations/20673990) PubMed: [20959462](http://www.uniprot.org/citations/20959462) PubMed: [22726440](http://www.uniprot.org/citations/22726440) PubMed: [24051492](http://www.uniprot.org/citations/24051492) PubMed: [9840937](http://www.uniprot.org/citations/9840937) PubMed: [24652652](http://www.uniprot.org/citations/24652652)). One of the activated genes is an inhibitor of cyclin-dependent kinases. Apoptosis induction seems to be mediated either by stimulation of BAX and FAS antigen expression, or by repression of Bcl-2 expression. Its pro-apoptotic activity is activated via its interaction with PPP1R13B/ASPP1 or TP53BP2/ASPP2 (PubMed: [12524540](http://www.uniprot.org/citations/12524540)). However, this activity is inhibited when the interaction with PPP1R13B/ASPP1 or TP53BP2/ASPP2 is displaced by PPP1R13L/iASPP (PubMed: [12524540](http://www.uniprot.org/citations/12524540)). In cooperation with mitochondrial PPIF is involved in activating oxidative stress-induced necrosis; the function is largely independent of transcription. Induces the transcription of long intergenic non-coding RNA p21 (lincRNA-p21) and lincRNA-Mkln1. LincRNA-p21 participates in TP53-dependent transcriptional repression leading to apoptosis and seems to have an effect on cell-cycle regulation. Implicated in Notch signaling cross-over. Prevents CDK7 kinase activity when associated to CAK complex in response to DNA damage, thus stopping cell cycle progression. Isoform 2 enhances the transactivation activity of isoform 1 from some but not all TP53-inducible promoters. Isoform 4 suppresses transactivation activity and impairs growth suppression mediated by isoform 1. Isoform 7 inhibits isoform 1-mediated apoptosis. Regulates the circadian clock by repressing CLOCK-BMAL1-mediated transcriptional activation of PER2 (PubMed: [24051492](http://www.uniprot.org/citations/24051492)).

Cellular Location

Cytoplasm. Nucleus. Nucleus, PML body Endoplasmic reticulum. Mitochondrion matrix. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome Note=Recruited into PML bodies together with CHEK2 (PubMed:12810724) Translocates to mitochondria upon oxidative stress (PubMed:22726440) Translocates to mitochondria in response to mitomycin C treatment (PubMed:27323408). Competitive inhibition of TP53 interaction with HSPA9/MOT-2 by UBXN2A results in increased protein abundance and subsequent translocation of TP53 to the nucleus (PubMed:24625977) [Isoform 2]: Nucleus. Cytoplasm. Note=Localized mainly in the nucleus with minor staining in the cytoplasm [Isoform 4]: Nucleus. Cytoplasm. Note=Predominantly nuclear but translocates to the cytoplasm following cell stress [Isoform 8]: Nucleus. Cytoplasm. Note=Localized in both nucleus and cytoplasm in most cells. In some cells, forms foci in the nucleus that are different from nucleoli

Tissue Location


Ubiquitous. Isoforms are expressed in a wide range of normal tissues but in a tissue-dependent manner. Isoform 2 is expressed in most normal tissues but is not detected in brain, lung, prostate, muscle, fetal brain, spinal cord and fetal liver. Isoform 3 is expressed in most normal tissues but is not detected in lung, spleen, testis, fetal brain, spinal cord and fetal liver. Isoform 7 is expressed in most normal tissues but is not detected in prostate, uterus, skeletal muscle and breast. Isoform 8 is detected only in colon, bone marrow, testis, fetal brain and intestine. Isoform 9 is expressed in most normal tissues but is not detected in brain, heart, lung, fetal liver, salivary gland, breast or intestine

p53 (C-terminus) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

p53 (C-terminus) Antibody - Images



Western blot detection of p53 in SW480, T47D and MDA-MB-468 cell lysates using p53 mouse mAb (1:1000 diluted). Predicted band size: 53KDa. Observed band size: 53KDa.

p53 (C-terminus) Antibody - Background

Acts as a tumor suppressor in many tumor types; induces growth arrest or apoptosis depending on the physiological circumstances and cell type. Involved in cell cycle regulation as a trans-activator that acts to negatively regulate cell division by controlling a set of genes required for this process. One of the activated genes is an inhibitor of cyclin-dependent kinases. Apoptosis induction seems to be mediated either by stimulation of BAX and FAS antigen expression, or by repression of Bcl-2 expression. In cooperation with mitochondrial PPIF is involved in activating oxidative stress-induced necrosis; the function is largely independent of transcription. Induces the transcription of long

intergenic non-coding RNA p21 (lincRNA-p21) and lincRNA- Mki1. LincRNA-p21 participates in TP53-dependent transcriptional repression leading to apoptosis and seem to have to effect on cell-cycle regulation. Implicated in Notch signaling cross-over. Prevents CDK7 kinase activity when associated to CAK complex in response to DNA damage, thus stopping cell cycle progression. Isoform 2 enhances the transactivation activity of isoform 1 from some but not all TP53-inducible promoters. Isoform 4 suppresses transactivation activity and impairs growth suppression mediated by isoform 1. Isoform 7 inhibits isoform 1-mediated apoptosis.

p53 (C-terminus) Antibody - References

Zakut-Houri R.,et al.EMBO J. 4:1251-1255(1985).
Lamb P.,et al.Mol. Cell. Biol. 6:1379-1385(1986).
Harlow E.,et al.Mol. Cell. Biol. 5:1601-1610(1985).
Harris N.,et al.Mol. Cell. Biol. 6:4650-4656(1986).
Buchman V.L.,et al.Gene 70:245-252(1988).