10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Hsp70 (C-terminus) Antibody
Purified Mouse Monoclonal Antibody (Mab)
Catalog # AP52677

Specification

Hsp70 (C-terminus) Antibody - Product Information

Application	**IP, WB**
Primary Accession	P08107
Reactivity	**Human**
Host	**Mouse**
Clonality	**Monoclonal**
Isotype	**IgG2b**
Calculated MW	**70 KDa**

Hsp70 (C-terminus) Antibody - Additional Information

Other Names
DAQB 147D11.1 001;FLJ54303;FLJ54370;FLJ54392;FLJ54408;FLJ75127;Heat shock 70 kDa protein 1;Heat shock 70 kDa protein 1/2;Heat shock 70 kDa protein 1A/1B;heat shock 70kDa protein 1A;Heat shock 70kDa protein 1B;Heat shock induced protein;heat shock protein 70;HSP70 1;HSP70 2;HSP70-1/HSP70-2;HSP70-1A;HSP70.1;HSP70.1/HSP70.2;HSP70I;HSP71_HUMAN;HSP72;HSPA1;HSPA1A;HSPA1B;XXbac BCX40G17.3 001.

Dilution
IP~~1:500
WB~~1:1000

Format
Purified mouse monoclonal in buffer containing 0.1M Tris-Glycine(pH 7.4,150 mM NaCl)with 0.09% (W/V) sodium azide,0.1mg/mlBSA and 50% glycerol.

Storage
Store at -20 °C.Stable for 12 months from date of receipt

Hsp70 (C-terminus) Antibody - Protein Information

Hsp70 (C-terminus) Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation


10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

- Flow Cytomety
- Cell Culture

Hsp70 (C-terminus) Antibody - Images



Immunoprecipitation analysis of Hela cell lysates using Hsp70 (C-terminus) mouse mAb.



Western blot detection of Hsp70(C-terminus) in Hela and HsHela cell lysates using Hsp70(C-terminus) mouse mAb (1:1000 diluted).Predicted band size: 70KDa.Observed band size: 70KDa.

Hsp70 (C-terminus) Antibody - Background

In cooperation with other chaperones, Hsp70s stabilize preexistent proteins against aggregation and mediate the folding of newly translated polypeptides in the cytosol as well as within organelles. These chaperones participate in all these processes through their ability to recognize nonnative conformations of other proteins. They bind extended peptide segments with a net hydrophobic character exposed by polypeptides during translation and membrane translocation, or following stress-induced damage. In case of rotavirus A infection, serves as a post-attachment receptor for the virus to facilitate entry into the cell.

Hsp70 (C-terminus) Antibody - References

Milner C.M.,et al.Immunogenetics 32:242-251(1990).
Hunt C.,et al.Proc. Natl. Acad. Sci. U.S.A. 82:6455-6459(1985).
Xie T.,et al.Genome Res. 13:2621-2636(2003).
Shiina S.,et al.Submitted (SEP-1999) to the EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).