10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Smad2 Antibody
Purified Mouse Monoclonal Antibody (Mab)
Catalog # AP52780

Specification

Smad2 Antibody - Product Information

Application	**WB, ICC**
Primary Accession	Q15796
Reactivity	**Human**
Host	**Mouse**
Clonality	**Monoclonal**
Isotype	**IgG1**
Calculated MW	**60 KDa**

Smad2 Antibody - Additional Information

Gene ID 4087

Other Names
hMAD 2;hMAD-2;hSMAD2;JV18 1;JV18;JV18;JV18-1;JV181;MAD;MAD;MAD homolog 2;MAD Related Protein 2;Mad-related protein 2;MADH2;MADR2;MGC22139;MGC34440;Mothers Against Decapentaplegic Homolog 2;Mothers Against Decapentaplegic Homolog 2;mothers against DPP homolog 2;OTTHUMP00000163489;Sma and Mad related protein 2;SMAD 2;SMAD;SMAD family member 2;SMAD, mothers against DPP homolog 2;SMAD2;SMAD2_HUMAN.

Dilution
WB~~1:500
ICC~~1:100

Format
ascites

Storage
Store at -20 °C.Stable for 12 months from date of receipt

Smad2 Antibody - Protein Information

Name SMAD2

Synonyms MADH2, MADR2

Function
Receptor-regulated SMAD (R-SMAD) that is an intracellular signal transducer and transcriptional modulator activated by TGF-beta (transforming growth factor) and activin type 1 receptor kinases. Binds the TRE element in the promoter region of many genes that are regulated by TGF-beta and, on formation of the SMAD2/SMAD4 complex, activates transcription. Promotes TGFB1-mediated transcription of odontoblastic differentiation genes in dental papilla cells (By similarity). Positively regulates PDPK1 kinase activity by stimulating its dissociation from the 14-3-3 protein YWHAQ

which acts as a negative regulator. May act as a tumor suppressor in colorectal carcinoma (PubMed:8752209).

Cellular Location
Cytoplasm. Nucleus. Note=Cytoplasmic and nuclear in the absence of TGF-beta. On TGF-beta stimulation, migrates to the nucleus when complexed with SMAD4 or with IPO7 (PubMed:9865696, PubMed:21145499). On dephosphorylation by phosphatase PPM1A, released from the SMAD2/SMAD4 complex, and exported out of the nucleus by interaction with RANBP1 (PubMed:16751101, PubMed:19289081). Localized mainly to the nucleus in the early stages of embryo development with expression becoming evident in the cytoplasm at the blastocyst and epiblast stages (By similarity). {ECO:0000250|UniProtKB:Q62432, ECO:0000269|PubMed:16751101, ECO:0000269|PubMed:19289081, ECO:0000269|PubMed:21145499, ECO:0000269|PubMed:9865696}


Tissue Location
Expressed at high levels in skeletal muscle, endothelial cells, heart and placenta.

Smad2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Smad2 Antibody - Images



Western blot detection of Smad2 in Hela,A431,Jurkat and K562 cell lysates using Smad2 mouse mAb (1:500 diluted).Predicted band size:60KDa.Observed band size:60KDa.

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999



Immunocytochemistry staining of HeLa cells fixed with 1% Paraformaldehyde and using Smad2 mouse mAb (dilution 1:100).

Smad2 Antibody - Background

Receptor-regulated SMAD (R-SMAD) that is an intracellular signal transducer and transcriptional modulator activated by TGF-beta (transforming growth factor) and activin type 1 receptor kinases. Binds the TRE element in the promoter region of many genes that are regulated by TGF-beta and, on formation of the SMAD2/SMAD4 complex, activates transcription. May act as a tumor suppressor in colorectal carcinoma. Positively regulates PDPK1 kinase activity by stimulating its dissociation from the 14-3-3 protein YWHAQ which acts as a negative regulator.

Smad2 Antibody - References

Riggins G.J.,et al.Nat. Genet. 13:347-349(1996).
Zhang Y.,et al.Nature 383:168-172(1996).
Eppert K.,et al.Cell 86:543-552(1996).
Liu F.,et al.Genes Dev. 11:3157-3167(1997).
Takenoshita S.,et al.Genomics 48:1-11(1998).