10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

CstF-64T Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP53338

Specification

CstF-64T Antibody - Product Information

Application	**WB**
Primary Accession	Q9H0L4
Reactivity	**Human**
Host	**Rabbit**
Clonality	**Polyclonal**
Calculated MW	**64 KDa**
Antigen Region	**91-140**

CstF-64T Antibody - Additional Information

Gene ID 23283

Dilution
WB~~ 1:1000

Format
Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.09% (W/V) sodium azide and 50% glycerol

Storage
Store at -20 °C.Stable for 12 months from date of receipt

CstF-64T Antibody - Protein Information

Name CSTF2T

Synonyms KIAA0689

Function
May play a significant role in AAUAAA-independent mRNA polyadenylation in germ cells. Directly involved in the binding to pre- mRNAs (By similarity).

Cellular Location
Nucleus.

CstF-64T Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

CstF-64T Antibody - Images



Anti-CstF-64T Antibody at 1:1000 dilution + Jurkat whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L),Peroxidase conjugated at 1/10000 dilution. Predicted band size : 64 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

CstF-64T Antibody - Background

May play a significant role in AAUAAA-independent mRNA polyadenylation in germ cells. Directly involved in the binding to pre-mRNAs (By similarity).

CstF-64T Antibody - References

Dass B.,et al.Genomics 80:509-514(2002).
Wiemann S.,et al.Genome Res. 11:422-435(2001).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Suzuki Y.,et al.Submitted (APR-2005) to the EMBL/GenBank/DDBJ databases.
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.